

Barresi and Gilbert, *Developmental Biology* 13e, Chapter 12 Literature Cited

Agassiz, A. and C. O. Whitman. 1884. On the development of some pelagic fish eggs—Preliminary notice. *Proc. Am. Acad. Arts Sci.* 20: 23–75.

Agathon, A., C. Thisse and B. Thisse. 2003. The molecular nature of the zebrafish tail organizer. *Nature* 424: 448–452.

Agius, E., M. Oelgeschläger, O. Wessely, C. Kemp and E. M. De Robertis. 2000. Endodermal Nodal-related signals and mesoderm induction in *Xenopus*. *Development* 127: 1173–1183.

Ancel, P. and P. Vintenberger. 1948. Recherches sur le determinisme de la symmetrie bilatérale dans l’oeuf des amphibiens. *Bull. Biol. Fr. Belg.* 31: 1–182.

Appel, T. A. 1987. *The Cuvier-Geoffroy Debate: French Biology in the Decades before Darwin*. Oxford University Press, New York.

Armon, R. 2012. Between biochemists and embryologists: The biochemical study of embryonic induction in the 1930s. *J. Hist. Biol.* 45: 65–108.

Bae, S, C. D. Reid and D. S. Kessler. 2011. Siamois and Twin are redundant and essential in formation of the Spemann organizer. *Dev. Biol.* 352: 367–381.

Beams, H. W. and R. G. Kessel. 1976. Cytokinesis: A comparative study of cytoplasmic division in animal cells. *Am. Sci.* 64: 279–290.

Beetschen, J. C. 2001. Amphibian gastrulation: History and evolution of a 125-year-old concept. *Int. J. Dev. Biol.* 45: 771–795.

Behrndt, M., G. Salbreux, P. Campinho, R. Hauschild, F. Oswald, J. Roensch, S. W. Grill and C. P. Heisenberg. 2012. Forces driving epithelial spreading in zebrafish gastrulation. *Science* 338: 257–260.

Bhattacharya, D., J. Zhong, S. Tavakoli, A. Kabla and P. Matsudaira. 2021. Strain maps characterize the symmetry of convergence and extension patterns during zebrafish gastrulation. *Sci. Rep.* 11: 19357.

Bier, E. and E. M. De Robertis. 2015. BMP gradients: A paradigm for morphogen-mediated developmental patterning. *Science* 348: 5838.

Bijtel, J. H. 1931. Über die Entwicklung des Schwanzes bei Amphibien. *Wilhelm Roux Arch. Entwicklungsmech. Org.* 125: 448–486.

Birsoy, B., M. Kofron, K. Schaible, C. Wylie and J. Heasman. 2006. Vg 1 is an essential signaling molecule in *Xenopus* development. *Development* 133: 15–20.

Blader, P. and U. Strähle 1998. Ethanol impairs migration of the prechordal plate in the zebrafish embryo. *Dev. Biol.* 201: 185–201.

Blitz, I. L. and K. W. Y. Cho. 1995. Anterior neurectoderm is progressively induced during gastrulation: The role of the *Xenopus* homeobox gene *orthodenticle*. *Development* 121: 993–1004.

Blum, M., A. Schweickert, P. Vick, C. V. Wright and M. V. Danilchik. 2014. Symmetry breakage in the vertebrate embryo: When does it happen and how does it work? *Dev. Biol.* 393: 109–123.

Blum, M., T. Beyer, T. Weber, P. Vick, P. Andre, E. Bitzer and A. Schweickert. 2009. *Xenopus*, an ideal model system to study vertebrate left-right asymmetry. *Dev. Dyn.* 238: 1215–1225.

Blythe, S. A., S. W. Cha, E. Tadjuidje, J. Heasman and P. S. Klein. 2010. Beta-Catenin primes organizer gene expression by recruiting a histone H3 arginine 8 methyltransferase, Prmt2. *Dev. Cell* 19: 220–231.

Boucaut, J.-C., T. D'Arribère, T. J. Poole, H. Aoyama, K. M. Yamada and J.-P. Thiery. 1984. Biologically active synthetic peptides as probes of embryonic development: A competitive peptide inhibition of fibronectin function inhibits gastrulation in amphibian embryos and neural crest cell migration in avian embryos. *J. Cell Biol.* 99: 1822–1830.

Bouwmeester, T., S.-H. Kim, Y. Sasai, B. Lu and E. M. De Robertis. 1996. Cerberus is a head-inducing secreted factor expressed in the anterior endoderm of Spemann's organizer. *Nature* 382: 595–601.

Bradbury, J. 2004. Small fish, big science. *PLOS Biology* 2: e148.

Brannon, M. and D. Kimelman. 1996. Activation of *siamois* by the Wnt pathway. *Dev. Biol.* 180: 344–347.

Braukmann, S. and S. F. Gilbert. 2005. Sucking in the gut: A history of early gastrulation research. In C. D. Stern (ed.), *Gastrulation: From Cells to Embryo*. Cold Spring Harbor Press, Cold Spring Harbor, NY, 1–20.

Bruce, A. E. E. 2016. Zebrafish epiboly: Spreading thin over the yolk. *Dev. Dyn.* 245: 244–258.

Caneparo, L., P. Pantazis, W. Dempsey, and S. E. Fraser. 2011. Intercellular bridges in vertebrate gastrulation. *PLOS ONE* 6: e20230.

Carmany-Rampey, A. and A. F. Schier. 2001. Single-cell internalization during zebrafish gastrulation. *Curr. Biol.* 11: 1261–1265.

Carnac, G., L. Kodjabachian, J. B. Gurdon and P. Lemaire. 1996. The homeobox gene *Siamois* is a target of the Wnt dorsalisation pathway and triggers organiser activity in the absence of mesoderm. *Development* 122: 3055–3065.

Carron, C. and D. L. Shi. 2016. Specification of anterioposterior axis by combinatorial signaling during *Xenopus* development. *Wiley Interdiscip. Rev. Dev. Biol.* 5: 150–168.

Carvalho, L. and C. P. Heisenberg. 2010. The yolk syncytial layer in early zebrafish development. *Trends Cell Biol.* 20: 586–592.

Cha, B. J. and D. L. Gard. 1999. XMAP230 is required for the organization of cortical microtubules and patterning of the dorsoventral axis in fertilized *Xenopus* eggs. *Dev. Biol.* 205: 275–286.

Chea, H. K., C. V. Wright and B. J. Swalla. 2005. Nodal signaling and the evolution of deuterostome gastrulation. *Dev. Dyn.* 234: 269–278.

Chien, Y. H., S. Srinivasan, R. Keller and C. Kintner. 2018. Mechanical strain determines cilia length, motility, and planar position in the left-right organizer. *Dev. Cell* 45: 316–330.e4.

Cho, K. W. Y. 2012. Enhancers. *Wiley Interdiscip. Rev. Dev. Biol.* 1: 469–478.

Chu, L. T., S. H. Fong, I. Kondrychyn, S. L. Loh, Z. Ye, and V. Korzh. 2012. Yolk syncytial layer formation is a failure of cytokinesis mediated by Rock1 function in the early zebrafish embryo. *Biol Open* 1: 747–753.

Cooper, M. S. and L. A. D'Amico. 1996. A cluster of noninvoluting endocytic cells at the margin of the zebrafish blastoderm marks the site of embryonic shield formation. *Dev. Biol.* 180: 184–198.

Cuykendall, T. N. and D. W. Houston. 2009. Vegetally localized *Xenopus* trim36 regulates cortical rotation and dorsal axis formation. *Development* 136: 3057–3065.

Darken, R. S., A. M. Scola, A. S. Rakeman, G. Das, M. Mlodzik and P. A. Wilson. 2002. The planar polarity gene *strabismus* regulates convergent extension movements in *Xenopus*. *EMBO J.* 21: 976–85.

Davidson, L. A., B. D. Dzamba, R. Keller and D. W. DeSimone. 2008. Live imaging of cell protrusive activity, and extracellular matrix assembly and remodeling during morphogenesis in the frog, *Xenopus laevis*. *Dev. Dyn.* 237: 2684–2692.

De Robertis, E. M. and J. Aréchaga (eds.). 2001. The Spemann-Mangold organizer: 75 years on. *Int. J. Dev. Biol.* 45 1–373.

De Robertis, E. M. and Y. Moriyama. 2016. The chordin morphogenetic pathway. *Curr. Top. Dev. Biol.* 116: 231–246.

De Robertis, E. M. and Y. Sasai. 1996. A common plan for dorsoventral patterning in Bilateria. *Nature* 380: 37–40.

De Robertis, E. M., J. Larraín, M. Oelgeschläger and O. Wessley. 2000. The establishment of Spemann's organizer and patterning of the vertebrate embryo. *Nat. Rev. Genet.* 1: 171–181.

Domingos, P. M., N. Itasaki, C. M. Jones, S. Mercurio, M. G. Sargent, J. C. Smith and R. Krumlauf. 2001. The Wnt/beta-catenin pathway posteriorizes neural tissue in *Xenopus* by an indirect mechanism requiring FGF signalling. *Dev. Biol.* 239: 148–160.

Dosch, R., D. S. Wagner, K. A. Mintzer, G. Runke, A. P. Wiemelt and M. C. Mullins. 2004. Maternal control of vertebrate development before the midblastula transition: Mutants from the zebrafish I. *Dev. Cell* 6: 771–780.

Dosch, R., V. Gawantka, H. Delius, C. Blumenstock and C. Niehrs. 1997. BMP-4 acts as a morphogen in dorsolateral mesoderm patterning in *Xenopus*. *Development* 124: 2325–2334.

Driever, W. 1995. Axis formation in zebrafish. *Curr. Opin. Genet. Dev.* 5: 610–618.

Driever, W. and 11 others. 1996. A genetic screen for mutations affecting development in zebrafish. *Development* 123: 37–46.

Du, S., B. W. Draper, M. Mione, C. B. Moens and A. Bruce. 2012. Differential regulation of epiboly initiation and progression by zebrafish Eomesodermin A. *Dev. Biol.* 362: 11–23.

Dumortier, J. G., S. Martin, D. Meyer, F. M. Rosa and N. B. David. 2012. Collective mesendoderm migration relies on an intrinsic directionality signal transmitted through cell contacts. *Proc. Natl. Acad. Sci. USA* 109: 16945–16950.

Durston, A. J., H. J. Jansen and S. A. Wacker. 2010a. Time-space translation: A developmental principle. *ScientificWorld* 10: 2207–2214.

Durston, A. J., H. J. Jansen and S. A. Wacker. 2010b. Time-space translation regulates trunk axial patterning in the early vertebrate embryo. *Genomics* 95: 250–255.

Elinson, R. P. and B. Rowning. 1988. A transient array of parallel microtubules in frog eggs: Potential tracks for a cytoplasmic rotation that specifies the dorso-ventral axis. *Dev. Biol.* 128: 185–197.

Elinson, R. P. and E. M. del Pino. 2012. Developmental diversity of amphibians. *Wiley Interdiscip. Rev. Dev. Biol.* 1: 345–369.

Engleka, M. J. and D. S. Kessler. 2001. Siamois cooperates with TGF β signals to induce the complete function of the Spemann-Mangold organizer. *Int. J. Dev. Biol.* 45: 241–250.

Essner, J. J., J. D. Amack, M. K. Nyholm, E. B. Harris and H. J. Yost. 2005. Kupffer's vesicle is a ciliated organ of asymmetry in the zebrafish embryo that initiates left-right development of the brain, heart and gut. *Development* 132: 1247–1260.

Essner, J. J., K. J. Vogan, M. K. Wagner, C. J. Tabin, H. J. Yost and M. Brueckner. 2002. Conserved function for embryonic nodal cilia. *Nature* 418: 37–38.

Ewald, A. J., H. McBride, M. Reddington, S. Fraser and R. Kerschmann. 2002. Surface imaging microscopy, an automated method for visualizing whole embryo samples in three dimensions at high resolution. *Dev. Dynam.* 225: 369–375.

Fan, M. J. and S. Y. Sokol. 1997. A role for Siamois in Spemann organizer formation. *Development* 124: 2581–2589.

Fauny, J. D., B. Thisse and C. Thisse. 2009. The entire zebrafish blastula-gastrula margin acts as an organizer dependent on the ratio of Nodal to BMP activity. *Development* 136: 3811–3819.

Fluck, R. A., K. L. Krok, B. A. Bast, S. E. Michaud and C. E. Kim. 1998. Gravity influences the position of the dorsoventral axis in medaka fish embryos (*Oryzias latipes*). *Dev. Growth Diff.* 40: 509–518.

Fukazawa, C., C. Santiago, K. M. Park, W. J. Deery, S. Gomez de la Torre Canny, C. K. Holterhoff and D. S. Wagner. 2010. *poky/chuk/ikk1* is required for differentiation of the zebrafish embryonic epidermis. *Dev. Biol.* 346: 272–283.

Fukuda, M., S. Takahashi, Y. Haramoto, Y. Onuma, Y. J. Kim, C. Y. Yeo, S. Ishiura and M. Asashima. 2010. Zygotic VegT is required for *Xenopus* paraxial mesoderm formation and is regulated by Nodal signaling and Eomesodermin. *Int. J. Dev. Biol.* 54: 81–92.

Funayama, N., F. Fagotto, P. McCrea and B. M. Gumbiner. 1995. Embryonic axis induction by the armadillo repeat domain of b-catenin: Evidence for intracellular signalling. *J. Cell Biol.* 128: 959–968.

Fürthauer, M., J. van Celst, C. Thisse and B. Thisse. 2004. FGF signaling controls the dorsoventral patterning of the zebrafish embryo. *Development* 131: 2853–2864.

Gawantka, V., H. Delius, K. Hirschfeld, C. Blumenstock and C. Niehrs. 1995. Antagonizing the Spemann organizer: Role of the homeobox gene *Xvent-1*. *EMBO J.* 14: 6268–6279.

Genikhovich, G., P. Fried, M. M. Prünster, J. B. Schinko, A. F. Gilles, D. Fredman, K. Meier, D. Iber and U. Technau. 2015. Axis patterning by BMPs: Cnidarian network reveals evolutionary

constraints. *Cell Rep.* 10 :1646–1654.

Gerhart, J. C., M. Danilchik, T. Doniach, S. Roberts, B. Rowning and R. Stewart. 1989. Cortical rotation of the *Xenopus* egg: Consequences for the anteroposterior pattern of embryonic dorsal development. *Development* 107: S37–S51.

Germain, S., M. Howell, G. M. Esslemont and C. S. Hill. 2000. Homeodomain and winged-helix transcription factors recruit activated Smads to distinct promoter elements via a common Smad interaction motif. *Genes Dev.* 14: 435–451.

Gilbert, S. F. and L. Saxén. 1993. Spemann's organizer: Models and molecules. *Mech. Dev.* 41: 73–89.

Glinka, A., W. Wu, D. Onichtchouk, C. Blumenstock and C. Niehrs. 1997. Head induction by simultaneous repression of BMP and Wnt signalling in *Xenopus*. *Nature* 389: 517–519.

Godsave, S. F. and J. M. W. Slack. 1989. Clonal analysis of mesoderm induction in *Xenopus laevis*. *Dev. Biol.* 134: 486–490.

Gont, L. K., H. Steinbeisser, B. Blumberg and E. M. De Robertis. 1993. Tail formation as a continuation of gastrulation: The multiple cell populations of the *Xenopus* tailbud derive from the late blastopore lip. *Development* 119: 991–1004.

Gonzales, A. P. and J. R. Yeh. 2014. Cas9-based genome editing in zebrafish. *Methods Enzymol.* 546: 377–413.

Goto, T., L. Davidson, M. Asashima and R. Keller. 2005. Planar cell polarity genes regulate polarized extracellular matrix deposition during frog gastrulation. *Curr. Biol.* 15: 787–793.

Granato, M. and C. Nüsslein-Volhard. 1996. Fishing for genes controlling development. *Curr. Opin. Genet. Dev.* 6: 461–468.

Gritsman, K., W. S. Talbot and A. F. Schier. 2000. Nodal signaling patterns the organizer. *Development* 127: 921–932.

Grunz, H. 1997. Neural induction in amphibians. *Curr. Topics Dev. Biol.* 35: 191–228.

Grunz, H. and L. Tacke. 1989. Neural differentiation of *Xenopus laevis* ectoderm takes place after disaggregation and delayed reaggregation without inducers. *Cell Diff. Dev.* 28: 211–217.

Guger, K. A. and B. M. Gumbiner. 1995. b-Catenin has wnt-like activity and mimics the Nieuwkoop signaling center in *Xenopus* dorsal-ventral patterning. *Dev. Biol.* 172: 115–125.

Haffter, P. and 16 others. 1996. The identification of genes with unique and essential functions in the development of the zebrafish, *Danio rerio*. *Development* 123: 1–36.

Hamburger, V. 1988. *The Heritage of Experimental Embryology: Hans Spemann and the organizer*. Oxford University Press, Oxford.

Hardin, J. D. and R. Keller. 1988. The behaviour and function of bottle cells during gastrulation of *Xenopus laevis*. *Development* 103: 211–230.

Hawley, S. H. B., K. Wünnenberg-Stapleton, C. Hashimoto, M. N. Laurent, T. Watabe, B. W. Blumberg and K. W. Y. Cho. 1995. Disruption of BMP signals in embryonic *Xenopus* ectoderm leads to direct neural induction. *Genes Dev.* 9: 2923–2935.

He, X., J.-P. Saint-Jeannet, J. R. Woodgett, H. E. Varmus and I. B. Dawid. 1995. Glycogen synthase kinase-3 and dorsoventral patterning in *Xenopus* embryos. *Nature* 374: 617–622.

Heasman, J., A. Crawford, K. Goldstone, P. Garner-Hamrick, B. Gumbiner, P. McCrea, C. Kinter, C. Y. Noro and C. Wylie. 1994a. Overexpression of cadherins and underexpression of b-catenin inhibit dorsal mesoderm induction in early *Xenopus* embryos. *Cell* 79: 791–803.

Heasman, J., D. Ginsberg, K. Goldstone, T. Pratt, C. Yoshidanaro and C. Wylie. 1994b. A functional test for maternally inherited cadherin in *Xenopus* shows its importance in cell adhesion at the blastula stage. *Development* 120: 49–57.

Helde, K. A., E. T. Wilson, C. J. Cretokos and D. J. Grunwald. 1994. Contribution of early cells to the fate map of the zebrafish gastrula. *Science* 265: 517–520.

Hemmati-Brivanlou A. and D. A. Melton. 1997. Vertebrate embryonic cells will become nerve cells unless told otherwise. *Cell* 88: 13–17.

Hemmati-Brivanlou, A. and D. A. Melton. 1992. A truncated activin receptor inhibits mesoderm induction and formation of axial structures in *Xenopus* embryos. *Nature* 359: 609–614.

Hemmati-Brivanlou, A. and D. A. Melton. 1994. Inhibition of activin signaling promotes neuralization in *Xenopus*. *Cell* 77: 273–281.

Hemmati-Brivanlou, A. and G. H. Thomsen. 1995. Ventral mesodermal patterning in *Xenopus* embryos: Expression patterns and activities of BMP-2 and BMP-4. *Dev. Genet.* 17: 78–89.

Holley, S. A., P. D. Jackson, Y. Sasai, B. Lu, E. M. De Robertis, F. M. Hoffmann and E. L. Ferguson. 1995. A conserved system for dorsal-ventral patterning in insects and vertebrates involving sog and chordin. *Nature* 376: 249–253.

Holtfreter, H. 1933. Die totale Exogastrulation, eine Selbststablösung des Ektoderms von Entomesoderm. Entwicklung und funktionelles Verhalten nervenloser Organe. *Arch. Entwickl. Mech. Org.* 129: 669–793.

Houliston, E. and R. P. Elinson. 1991. Evidence for the involvement of microtubules, endoplasmic reticulum, and kinesin in cortical rotation of fertilized frog eggs. *J. Cell Biol.* 114: 1017–1028.

Hurtado, C. and E. M. De Robertis. 2007. Neural induction in the absence of organizer in salamanders is mediated by MAPK. *Dev. Biol.* 307: 282–289.

Ibrahim, H. and R. Winklbauer. 2001. Mechanisms of mesendoderm internalization in *Xenopus* gastrula: Lessons from the ventral side. *Dev. Biol.* 240: 108–122.

Iemura, S.-I., T. S. Yamamoto, C. Takagi, H. Uchiyama, T. Natsume, S. Shimasaki, H. Sugino and N. Ueno. 1998. Direct binding of follistatin to a complex of bone morphogenetic protein and its receptor inhibits ventral and epidermal cell fates in early *Xenopus* embryo. *Proc. Natl. Acad. Sci. USA* 95: 9337–9342.

Jessen, J. R., J. Topczewski, S. Bingham, D. S. Sepich, F. Marlow, A. Chandrasekhar and L. Solnica-Krezel. 2002. Zebrafish trilobite identifies new roles for Strabismus in gastrulation and neuronal movements. *Nat. Cell Biol.* 4: 610–615.

Kane, D. A. and C. B. Kimmel. 1993. The zebrafish midblastula transition. *Development* 119: 447–456.

Kane, D. A., K. N. McFarland and R. M. Warga. 2005. Mutations in half baked/E-cadherin block cell behaviors that are necessary for teleost epiboly. *Development* 132: 1105–1116.

Keller, P. J., A. D. Schmidt, J. Wittbrodt and E. H. K. Stelzer. 2008. Reconstruction of zebrafish early embryonic development by scanned light sheet microscopy. *Science* 322: 1065–1069.

Keller, R. E. 1975. Vital dye mapping of the gastrula and neurula of *Xenopus laevis*. I. Prospective areas and morphogenetic movements of the superficial layer. *Dev. Biol.* 42: 222–241.

Keller, R. E. 1976. Vital dye mapping of the gastrula and neurula of *Xenopus laevis*. II. Prospective areas and morphogenetic movements of the deep layer. *Dev. Biol.* 51: 118–137.

Keller, R. E. 1986. The cellular basis of amphibian gastrulation. In L. Browder (ed.), *Developmental Biology: A Comprehensive Synthesis*, Vol. 2. Plenum, New York, 241–327.

Keller, R. E. and G. C. Schoenwolf. 1977. An SEM study of cellular morphology, contact, and arrangement as related to gastrulation in *Xenopus laevis*. *Wilhelm Roux Arch. Dev. Biol.* 182: 165–186.

Keller, R. and M. Danilchik. 1988. Regional expression, pattern and timing of convergence and extension during gastrulation of *Xenopus laevis*. *Development* 103: 193–209.

Keller, R. and A. Sutherland. 2020. Convergent extension in the amphibian, *Xenopus laevis*. *Curr. Top. Dev. Biol.* 136: 271–317.

Keller, R., L. A. Davidson and D. R. Shook. 2003. How we are shaped: The biomechanics of gastrulation. *Differentiation* 71: 171–205.

Kelly, G. M., D. F. Erezilmaz and R. T. Moon. 1995. Induction of a secondary embryonic axis in zebrafish occurs following the overexpression of b-catenin. *Mech. Dev.* 53: 261–273.

Kessler, D. S. 1997. Siamois is required for formation of Spemann's organizer. *Proc. Natl. Acad. Sci. USA* 94: 13017–13022.

Khokha, M. K., J. Yeh, T. C. Grammer and R. M. Harland. 2005. Depletion of three BMP antagonists from Spemann's organizer leads to catastrophic loss of dorsal structures. *Dev. Cell* 8: 401–411.

Kimmel, C. B. and R. D. Law. 1985. Cell lineage of zebrafish blastomeres. II. Formation of the yolk syncytial layer. *Dev. Biol.* 108: 86–93.

Kimmel, C. B. and R. M. Warga. 1987. Indeterminate cell lineage of the zebrafish embryo. *Dev. Biol.* 124: 269–280.

Kimmel, C. B., W. W. Ballard, S. R. Kimmel, B. Ullmann and T. F. Schilling. 1995. Stages of embryonic development in the zebrafish. *Dev. Dyn.* 203: 253–310.

Kishimoto, Y., S. Koshita, M. Furutani-Seiki and H. Kondoh. 2004. Zebrafish maternal-effect mutations causing cytokinesis defect without affecting mitosis or equatorial *vasa* deposition. *Mech. Dev.* 121: 79–89.

Klein, S. L. and S. A. Moody. 2015. Early neural ectodermal genes are activated by Siamois and Twin during blastula stages. *Genesis* 53: 308–320.

Kofron, M., T. Demel, J. Xanthos, J. Lohr, B. Sun, H. Sive, S. Osada, C. Wright, C. Wylie and J. Heasman. 1999. Mesoderm induction in *Xenopus* is a zygotic event regulated by maternal VegT via TGF β growth factors. *Development* 126: 5759–5770.

Kornikova, E. S., E. G. Korvin-Pavlovskaya and L. V. Beloussov. 2009. Relocations of cell convergence sites and formation of pharyngula-like shapes in mechanically relaxed *Xenopus* embryos. *Dev. Genes Evol.* 219: 1–10.

Koshida, S., M. Shinya, T. Mizuno, A. Kuroiwa and H. Takeda. 1998. Initial anteroposterior pattern of zebrafish central nervous system is determined by differential competence of the epiblast. *Development* 125: 1957–1966.

Ku, M. and D. A. Melton. 1993. *Xwnt-11*: A maternally expressed *Xenopus wnt* gene. *Development* 119: 1161–1173.

Kubo, H., K. Shiga, Y. Harada and Y. Iwao. 2010. Analysis of a sperm surface molecule that binds to a vitelline envelope component of *Xenopus laevis* eggs. *Mol. Reprod. Dev.* 77: 728–735.

Lamb, T. M. and 7 others. 1993. Neural induction by the secreted polypeptide noggin. *Science* 262: 713–718.

- Landström, U. and S. Løvtrup. 1979. Fate maps and cell differentiation in the amphibian embryo: An experimental study. *J. Embryol. Exp. Morphol.* 54: 113–130.
- Lane, M. C. and W. C. Smith. 1999. The origins of primitive blood in *Xenopus*: Implications for axial patterning. *Development* 126: 423–434.
- Langdon, Y. G. and M. C. Mullins. 2011. Maternal and zygotic control of zebrafish dorsoventral axial patterning. *Annu. Rev. Genet.* 45: 357–377.
- Langeland, J. and C. B. Kimmel. 1997. The embryology of fish. In S. F. Gilbert and A. M. Raunio (eds.), *Embryology: Constructing the Organism*. Sinauer Associates, Sunderland, MA, 383–407.
- Larabell, C. A. and 7 others. 1997. Establishment of the dorsal-ventral axis in *Xenopus* embryos is presaged by early asymmetries in beta-catenin which are modulated by the Wnt signaling pathway. *J. Cell Biol.* 136: 1123–1136.
- Laurent, M. N., I. L. Blitz, C. Hashimoto, U. Rothbächer and K. W.-Y. Cho. 1997. The *Xenopus* homeobox gene *twin* mediates Wnt induction of *gooseoid* in establishment of Spemann's organizer. *Development* 124: 4905–4916.
- Lee, K. W., S. E. Webb and A. L. Miller. 2003. Ca^{2+} released via IP_3 receptors is required for furrow deepening during cytokinesis in zebrafish embryos. *Int. J. Dev. Biol.* 47: 411–421.
- Lemaire, P., N. Garrett and J. B. Gurdon. 1995. Expression cloning of *Siamois*, a *Xenopus* homeobox gene expressed in dorsal-vegetal cells of blastulae and able to induce a complete secondary axis. *Cell* 81: 85–94.
- Lepage, S. E. and A. E. Bruce. 2010. Zebrafish epiboly: Mechanics and mechanisms. *Int. J. Dev. Biol.* 54: 1213–1228.
- Leslie, J. D. and R. Mayor. 2013. Complement in animal development: Unexpected roles of a highly conserved pathway. *Semin. Immunol.* 25: 39–46.
- Leung, C., S. E. Webb and A. L. Miller. 1998. Calcium transients accompany ooplasmic segregation in zebrafish embryos. *Dev. Growth Diff.* 40: 313–326.

Leung, C., S. E. Webb and A. L. Miller. 2000. On the mechanism of ooplasmic segregation in single-cell zebrafish embryos. *Dev. Growth Diff.* 42: 29–40.

Lobikin, M. G. Wang, J. Xu, Y. W. Hsieh, C. F. Chuang, J. M. Lemire and M. Levin. 2012. Early, nonciliary role for microtubule proteins in left-right patterning is conserved across kingdoms. *Proc. Natl. Acad. Sci. USA* 109: 12586–12591.

Long, S., N. Ahmad and M. Rebagliati. 2003. The zebrafish nodal-related gene *southpaw* is required for visceral and diencephalic left-right asymmetry. *Development* 130: 2303–2316.

Lu, F. I., C. Thisse and B. Thisse. 2011. Identification and mechanism of regulation of the zebrafish dorsal determinant. *Proc. Natl. Acad. Sci. USA* 108: 15876–15880.

Manes, M. E. and R. P. Elinson. 1980. Ultraviolet light inhibits gray crescent formation in the frog egg. *Wilhelm Roux Arch. Dev. Biol.* 189: 73–77.

Mangold, O. 1933. Über die Induktionsfähigkeit der verschiedenen Bezirke der Neurula von Urodelen. *Naturwissenschaften* 21: 761–766.

Marsden, M. and D. W. DeSimone. 2001. Regulation of cell polarity, radial intercalation, and epiboly in *Xenopus*: Novel roles for integrin and fibronectin. *Development* 128: 3635–3647.

McFarland, K. N., R. M. Warga and D. A. Kane. 2005. Genetic locus *half baked* is necessary for morphogenesis of the ectoderm. *Dev. Dyn.* 233: 390–406.

McMahon, A. P. and R. T. Moon. 1989. Ectopic expression of the proto-oncogene *int-1* in *Xenopus* embryos leads to duplication of the embryonic axis. *Cell* 58: 1075–1084.

Miller, J. R., B. A. Rowling, C. A. Larabell, J. A. Yang-Snyder, R. L. Bates and R. T. Moon. 1999. Establishment of the dorsal-ventral axis in *Xenopus* embryos coincides with the dorsal enrichment of Dishevelled that is dependent on cortical rotation. *J. Cell Biol.* 146: 427–437.

Molenaar, M. and 8 others. 1996. XTcf-3 transcription factor mediates beta-catenin-induced axis formation in *Xenopus* embryos. *Cell* 86: 391–399.

Moon, R. T. and D. Kimelman. 1998. From cortical rotation to organizer gene expression: Toward a molecular explanation of axis specification in *Xenopus*. *BioEssays* 20: 536–545.

Moosmann, J., A. Ershov, V. Altapova, T. Baumbach, M. S. Prasad, C. LaBonne, X. Xiao, J. Kashef and R. Hofmann. 2013. X-ray phase-contrast in vivo microtomography probes new aspects of *Xenopus* gastrulation. *Nature* 497: 574–377.

Nagai, K. and 7 others. 2009. The sperm-surface glycoprotein SGP is necessary for fertilization in the frog *Xenopus laevis*. *Dev. Growth Differ.* 51: 499–510.

Nakamura, O. and H. Takasaki. 1970. Further studies on the differentiation capacity of the dorsal marginal zone in the morula of *Triturus pyrrhogaster*. *Proc. Jpn. Acad.* 46: 700–705.

Newman, C. S. and P. A. Krieg. 1999. Specification and differentiation of the heart in amphibia. In S. A. Moody (ed.), *Cell Lineage and Fate Determination*. Academic Press, New York, 341–351.

Newport, J. W. and M. W. Kirschner. 1982a. A major developmental transition in early *Xenopus* embryos: I. Characterization and timing of cellular changes at midblastula stage. *Cell* 30: 675–686.

Newport, J. W. and M. W. Kirschner. 1982b. A major developmental transition in early *Xenopus* embryos. II. Control of the onset of transcription. *Cell* 30: 687–696.

Niehrs, C. 2004. Regionally specific induction by the Spemann-Mangold organizer. *Nat. Rev. Genet.* 5: 425–434.

Nieuwkoop, P. D. 1969. The formation of the mesoderm in urodele amphibians. I. Induction by the endoderm. *Wilhelm Roux Arch. Entwicklungsmech. Org.* 162: 341–373.

Nieuwkoop, P. D. 1973. The “organisation center” of the amphibian embryo: Its origin, spatial organisation and morphogenetic action. *Adv. Morphogenet.* 10: 1–39.

Nieuwkoop, P. D. 1977. Origin and establishment of embryonic polar axes in amphibian development. *Curr. Top. Dev. Biol.* 11: 115–132.

Nieuwkoop, P. D. and P. A. Florschütz. 1950. Quelques caractères spéciaux de la gastrulation et de la neurulation de l’oeuf de *Xenopus laevis*, Daud. et de quelques autres anoures. *Arch. Biol.* 61: 113–150.

Northrop, J., A. Woods, R. Seger, A. Suzuki, N. Ueno, E. Krebs and D. Kimelman. 1995. BMP-4 regulates the dorsal-ventral differences in FGF/MAPK-mediated mesoderm induction in

Xenopus. *Dev Biol.* 172: 242–252.

Oelgeschläger, M., H. Kuroda, B. Reversade and E. M. De Robertis. 2003. Chordin is required for the Spemann organizer transplantation phenomenon in *Xenopus* embryos. *Dev. Cell* 4: 219–230.

Okada, Y., S. Tanaka, Y. Tanaka, J.-C. Belmonte and N. Hirokawa. 2005. Mechanism of nodal flow: A conserved breaking event in left-right axis determination. *Cell* 121: 633–644.

Onuma, Y., S. Takahashi, C. Yokota and M. Asashima. 2002. Multiple nodal-related genes act coordinately in *Xenopus* embryogenesis. *Dev. Biol.* 241: 94–105.

Oppenheimer, J. M. 1936. Transplantation experiments on developing teleosts (*Fundulus* and *Perca*). *J. Exp. Zool.* 72: 409–437.

Ossipova, O., C. W. Chu, J. Fillatre, B. K. Brott, K. Itoh, and S. Y. Sokol. 2015. The involvement of PCP proteins in radial cell intercalations during *Xenopus* embryonic development. *Dev. Biol.* 408: 316–327.

Papan, C., B. Boulat, S. S. Velan, S. E. Fraser and R. E. Jacobs. 2007a. Formation of the dorsal marginal zone in *Xenopus laevis* analyzed by time-lapse microscopic magnetic resonance imaging. *Dev. Biol.* 305: 161–171.

Papan, C., B. Boulat, S. Velan, S. E. Fraser, and R. E. Jacobs. 2007b. Two-dimensional and three-dimensional time-lapse microscopic magnetic resonance imaging of *Xenopus* gastrulation movements using intrinsic tissue-specific contrast. *Dev. Dyn.* 236: 494–501.

Pera, E. M., H. Acosta, N. Gougnard, M. Climent and I. Arregi. 2014. Active signals, gradient formation, and regional specificity in neural induction. *Exp. Cell Res.* 321: 25–31.

Pera, E. M., O. Wessely, S.-S. Li and E. M. De Robertis. 2001. Neural and head induction by insulin-like growth factor signals. *Dev. Cell* 1: 655–665.

Petersen, C. P. and P. W. Reddien. 2009. Wnt signaling and the polarity of the primary body axis. *Cell* 139: 1056–1068.

Pézeron, G., P. Mourrain, S. Courty, J. Ghislain, T. S. Becker, F. M. Rosa and N. B. David. 2008. Live analysis of endodermal layer formation identifies random walk as a novel gastrulation

movement. *Curr. Biol.* 18: 276–281.

Pfister, K., D. R. Shook, C. Chang, R. Keller and P. Skoglund. 2016. Molecular model for force production and transmission during vertebrate gastrulation. *Development* 143: 715–727.

Piccolo, S., E. Agius, L. Leyns, S. Bhattacharyya, H. Grunz, T. Bouwmeester and E. M. DeRobertis. 1999. The head inducer Cerberus is a multifunctional antagonist of Nodal, BMP, and Wnt signals. *Nature* 397: 707–710.

Piccolo, S., Y. Sasai, B. Lu and E. M. De Robertis. 1996. Dorsoventral patterning in *Xenopus*: Inhibition of ventral signals by direct binding of chordin to BMP-4. *Cell* 86: 589–598.

Pierce, S. B. and D. Kimelman. 1995. Regulation of Spemann organizer formation by the intracellular kinase Xgsk-3. *Development* 121: 755–765.

Rankin, S. A., J. Kormish, M. Kofron, A. Jegga and A. M. Zorn. 2011. A gene regulatory network controlling *hhx* transcription in the anterior endoderm of the organizer. *Dev. Biol.* 351: 297–310.

Rebagliati, M. R., R. Toyama, C. Fricke, P. Haffter and I. B. Dawid. 1998. Zebrafish nodal-related genes are implicated in axial patterning and establishing left-right asymmetry. *Dev. Biol.* 199: 261–272.

Rex, M., E. Hilton and R. Old. 2002. Multiple interactions between maternally-activated signaling pathway control *Xenopus* nodal-related genes. *Int. J. Dev. Biol.* 46: 217–226.

Richard-Parpaillon, L., C. Héligon, F. Chesnel, D. Boujard and A. Philpott. 2002. The IGF pathway regulates head formation by inhibiting Wnt signaling in *Xenopus*. *Dev. Biol.* 244: 407–417.

Rogers, C. D., N. Harafuji, T. Archer, D. D. Cunningham and E. S. Casey. 2009a. *Xenopus* Sox3 activates *sox2* and *geminin* and indirectly represses *Xvent2* expression to induce neural progenitor formation at the expense of non-neural ectodermal derivatives. *Mech. Dev.* 126: 42–55.

Rogers, C. D., S. A. Moody and E. S. Casey. 2009b. Neural induction and factors that stabilize a neural fate. *Birth Defects Res. C: Embryol. Today* 87: 249–262.

Rogers, C. D., T. C. Archer, D. D. Cunningham, T. C. Grammer and E. M. Casey. 2008. Sox3 expression is maintained by FGF signaling and restricted to the neural plate by Vent proteins in the *Xenopus* embryo. *Dev. Biol.* 313: 307–319.

Rogers, K. W. and P. Müller. 2018. Nodal and BMP dispersal during early zebrafish development. *Dev. Biol.* 447: 14–23.

Romeo, N., A. Hastewell, A. Mietke and J. Dunkel 2021. Learning developmental mode dynamics from single-cell trajectories. *Elife* 10:e68679.

Roux, W. 1887. Beiträge zur Entwicklungsmechanik des Embryo. *Arch. Mikrosk. Anat.* 29: 157–212.

Royer, L. A., W. C. Lemon, R. K. Chhetri, Y. Wan, M. Coleman, E. W. Myers and P. J. Keller. 2016. Adaptive light-sheet microscopy for long-term, high-resolution imaging in living organisms. *Nat. Biotechnol.* 34: 1267–1278.

Rozario, T., B. Dzamba, G. F. Weber, L. A. Davidson and D. W. DeSimone. 2009. The physical state of fibronectin matrix differentially regulates morphogenetic movements in vivo. *Dev. Biol.* 327: 386–398.

Ryan, A. K. and 14 others. 1998. Pitx2 determines left-right asymmetry of internal organs in vertebrates. *Nature* 394: 545–551.

Saka, Y. and J. C. Smith. 2001. Spatial and temporal patterns of cell division during early *Xenopus* embryogenesis. *Dev. Biol.* 229: 307–318.

Sampath, K., A. L. Rubinstein, A. M. Cheng, J. O. Liang, K. Fekany, L. Solnica-Krezel, V. Korzh, M. E. Halpern and C. V. Wright. 1998. Induction of the zebrafish ventral brain and floorplate requires cyclops/nodal signalling. *Nature* 395: 185–189.

Sander, K. and P. E. Faessler. 2001. Introducing the Spemann-Mangold organizer: Experiments and insights that generated a key concept in developmental biology. *Int. J. Dev. Biol.* 45: 1–11.

Sasai, Y., B. Lu, H. Steinbeisser, D. Geissert, L. K. Gont and E. M. De Robertis. 1994. *Xenopus* chordin: A novel dorsalizing factor activated by organizer-specific homeobox genes. *Cell* 79: 779–790.

Sasai, Y., B. Lu, S. Piccolo and E. M. de Robertis. 1996. Endoderm induction by the organizer-secreted factors chordin and noggin in *Xenopus* animal caps. *EMBO J.* 15: 4547–4555.

Sato, S. M. and T. D. Sargent. 1989. Development of neural inducing capacity in dissociated *Xenopus* embryos. *Dev. Biol.* 134: 263–266.

Saxén, L. 1961. Transfilter neural induction of amphibian ectoderm. *Dev. Biol.* 3: 140–152.

Saxén, L. 2001. Spemann's heritage in Finnish developmental biology. *Int. J. Dev. Biol.* 45: 51–55.

Saxén, L. and S. Toivonen. 1962. *Embryonic Induction*. Prentice-Hall, Englewood Cliffs, NJ.

Schier, A. F. 2001. Axis formation and patterning in zebrafish. *Curr. Opin. Genet. Dev.* 11: 393–404.

Schier, A. F. and W. S. Talbot. 1998. The zebrafish organizer. *Curr. Opin. Genet. Dev.* 8: 464–471.

Schier, A. F., S. C. Neuhauss, K. A. Held, W. S. Talbot and W. Driever. 1997 The *one eyed pinhead* gene functions in mesoderm and endoderm formation in zebrafish and interacts with *no tail*. *Development* 124: 327–342.

Schmitz, B. and J. A. Campos-Ortega. 1994. Dorso-ventral polarity of the zebrafish embryo is distinguishable prior to the onset of gastrulation. *Wilhelm Roux Arch. Dev. Biol.* 203: 374–380.

Schneider, S., H. Steinbeisser, R. M. Warga and P. Hausen. 1996. Beta-catenin translocation into nuclei demarcates the dorsalizing centers in frog and fish embryos. *Mech. Dev.* 57: 191–198.

Schroeder, K. E., M. L. Condic, L. M. Eisenberg and H. J. Yost. 1999. Spatially regulated translation in embryos: Asymmetric expression of maternal Wnt-11 along the dorsal-ventral axis in *Xenopus*. *Dev. Biol.* 214: 288–297.

Schweickert, A., T. Weber, T. Beyer, P. Vick, S. Bogusch, K. Feistel and M. Blum. 2007. Cilia-driven leftward flow determines laterality in *Xenopus*. *Curr. Biol.* 17: 60–66.

- Shah, G. and 8 others. 2019. Multi-scale imaging and analysis identify pan-embryo cell dynamics of germlayer formation in zebrafish. *Nat. Commun.* 10: 5753.
- Shimizu, T., T. Yabe, O. Muraoka, S. Yonemura, S. Aramaki, K. Hatta, Y. K. Bae, H. Nojima and M. Hibi. 2005a. E-cadherin is required for gastrulation cell movements in zebrafish. *Mech. Dev.* 122: 747–763.
- Shimizu, T., Y. K. Bae, O. Muraoka and M. Hibi. 2005b. Interaction of Wnt and caudal-related genes in zebrafish posterior body formation. *Dev. Biol.* 279: 125–141.
- Shindo, A. 2018. Models of convergent extension during morphogenesis. *Wiley Interdiscip. Rev. Dev. Biol.* 7: e293.
- Shindo, A. and J. B. Wallingford. 2014. PCP and septins compartmentalize cortical actomyosin to direct collective cell movement. *Science* 343: 649–652.
- Shindo, A., Y. Inoue, M. Kinoshita and J. B. Wallingford. 2019. PCP-dependent transcellular regulation of actomyosin oscillation facilitates convergent extension of vertebrate tissue. *Dev. Biol.* 446: 159–167.
- Shinya, M., M. Furutani-Seiki, A. Kuroiwa and H. Takeda. 1999. Mosaic analysis with *oep* mutant reveals a repressive interaction between floor-plate and non-floor-plate mutant cells in the zebrafish neural tube. *Dev. Growth Diff.* 41: 135–142.
- Shook, D. R., E. M. Kasproicz, L. A. Davidson and R. Keller. 2018. Large, long-range tensile forces drive convergence during *Xenopus* blastopore closure and body axis elongation. *Elife* 7: e26944.
- Siddiqui, M., H. Sheikh, C. Tran, A. E. Bruce. 2010. The tight junction component Claudin E is required for zebrafish epiboly. *Dev. Dyn.* 239: 715–722.
- Silva, A. C., M. Filipe, K.-M. Kuerner, H. Steinbeisser and J. A. Belo. 2003. Endogenous Cerberus activity is required for anterior head specification in *Xenopus*. *Development* 130: 4943–4953.
- Skirkanich, J., G. Luxardi, J. Yang, L. Kodjabachian and P. S. Klein. 2011. An essential role for transcription before the MBT in *Xenopus laevis*. *Dev. Biol.* 357: 478–491.

- Smith, J. C. and J. M. W. Slack. 1983. Dorsalization and neural induction: Properties of the organizer in *Xenopus laevis*. *J. Embryol. Exp. Morphol.* 78: 299–317.
- Smith, W. C. and R. M. Harland. 1991. Injected Xwnt-8 RNA acts early in *Xenopus* embryos to promote formation of a vegetal dorsalizing center. *Cell* 67: 753–765.
- Smith, W. C. and R. M. Harland. 1992. Expression cloning of noggin, a new dorsalizing factor localized to the Spemann organizer in *Xenopus* embryos. *Cell* 70: 829–840.
- Smith, W. C., A. K. Knecht, M. Wu and R. M. Harland. 1993. Secreted noggin mimics the Spemann organizer in dorsalizing *Xenopus* mesoderm. *Nature* 361: 547–549.
- Smithers, L. E., C. M. Jones. 2002. *Xhex*-expressing endodermal tissues are essential for anterior patterning in *Xenopus*. *Mech. Dev.* 119: 191–200.
- Sokol, S., J. L. Christian, R. T. Moon and D. A. Melton. 1991. Injected Wnt RNA induces a complete body axis in *Xenopus* embryos. *Cell* 67: 741–752.
- Solnica-Krezel, L. and W. Driever. 1994. Microtubule arrays of the zebrafish yolk cell: Organization and function during epiboly. *Development* 120: 2443–2455.
- Solnica-Krezel, L. and W. Driever. 2001. The role of the homeodomain protein Bozozok in zebrafish axis formation. *Int. J. Dev. Biol.* 45: 299–310.
- Sonavane, P. R., C. Wang, B. Dzamba, G. F. Weber, A. Periasamy and D. W. DeSimone. 2017. Mechanical and signaling roles for keratin intermediate filaments in the assembly and morphogenesis of *Xenopus* mesendoderm tissue at gastrulation. *Development* 144: 4363–4376.
- Stancheva, I., O. El-Maarri, J. Walter, A. Niveleau and R. R. Meehan. 2002. DNA methylation at promoter regions regulates the timing of gene activation in *Xenopus laevis* embryos. *Dev. Biol.* 243: 155–165.
- Steinbeisser, H., A. Fainsod, C. Niehrs, Y. Sasai and E. M. De Robertis. 1995. The role of gsc and BMP-4 in dorsal-ventral patterning of the marginal zone in *Xenopus*: A loss-of-function study using antisense RNA. *EMBO J.* 14: 5230–5243.

Strahle, U. and S. Jesuthasan. 1993. Ultraviolet irradiation impairs epiboly in zebrafish embryos: Evidence for a microtubule-dependent mechanism of epiboly. *Development* 119: 909–919.

Szabó, A., I. Cobo, S. Omara, S. McLachlan, R. Keller and R. Mayor. 2016. The molecular basis of radial intercalation during tissue spreading in early development. *Dev. Cell* 37: 213–225.

Tao, Q. and 9 others. 2005. Maternal *wnt11* activates the canonical Wnt signaling pathway required for axis formation in *Xenopus* embryos. *Cell* 120: 857–871.

Taverner, N. V. and 8 others. 2005. Microarray-based identification of VegT targets in *Xenopus*. *Mech. Dev.* 122: 333–354.

Thisse, B. and C. Thisse. 2015. Formation of the vertebrate embryo: Moving beyond the Spemann organizer. *Semin. Cell Dev. Biol.* 42: 94–102.

Thisse, B., C. V. Wright and C. Thisse. 2000. Activin- and Nodal-related factors control antero-posterior patterning of the zebrafish embryo. *Nature* 403: 425–428.

Toivonen, S. and J. Wartiovaara. 1976. Mechanism of cell interaction during primary induction studied in transfilter experiments. *Differentiation* 5: 61–66.

Toivonen, S. and L. Saxén 1968. Morphogenetic interaction of presumptive neural and mesodermal cells mixed in different ratios. *Science* 159: 539–540.

Toivonen, S. and L. Saxén. 1955. The simultaneous inducing action of liver and bone marrow of the guinea pig in implantation and explanation experiments with embryos of *Triturus*. *Exp. Cell Res.* 3: 346–357.

Toivonen, S., D. Tarin, L. Saxén, P. J. Tarin and J. Wartiovaara. 1975. Transfilter studies on neural induction in the newt. *Differentiation* 4: 1–7.

Trinkaus, J. P. 1984. Mechanisms of *Fundulus* epiboly: A current view. *Am. Zool.* 24: 673–688.

Trinkaus, J. P. 1992. The midblastula transition, the YSL transition, and the onset of gastrulation in *Fundulus*. *Development* 1992: 75–80.

- Trinkaus, J. P. 1993. The yolk syncitial layer of *Fundulus*: Its origin and history and its significance for early embryogenesis. *J. Exp. Zool.* 265: 258–284.
- Tsang, M., S. Maegawa, A. Kiang, R. Habas, E. Weinberg and I. B. Dawid. 2004. A role for MKP3 in axial patterning of the zebrafish embryo. *Development* 131: 2769–2779.
- Tuazon, F. B. and M. C. Mullins. 2015. Temporally coordinated signals progressively pattern the anteroposterior and dorsoventral body axes. *Semin. Cell Dev. Biol.* 42: 118–133.
- Tucker, A. S. and J. M. Slack. 1995. Tail bud determination in the vertebrate embryo. *Curr. Biol.* 5: 807–813.
- Twitty, V. C. 1966. *Of Scientists and Salamanders*. Freeman, San Francisco, CA.
- Valles, J. M., Jr., S. R. Wasserman, C. Schweidenback, J. Edwardson, J. M. Denegre and K. L. Mowry. 2002. Processes that occur before second cleavage determine third cleavage orientation in *Xenopus*. *Exp. Cell Res.* 274: 112–118.
- Varshney, G. K., R. Sood and S. M. Burgess. 2015. Understanding and editing the zebrafish genome. *Adv. Genet.* 92: 1–52.
- Vervenne, H. B., K. R. Crombez, K. Lambaerts, L. Carvalho, M. Köppen, C. P. Heisenberg, W. J. Van de Ven and M. M. Petit. 2008. Lpp is involved in Wnt/PCP signaling and acts together with Scrib to mediate convergence and extension movements during zebrafish gastrulation. *Dev. Biol.* 320: 267–277.
- Vincent, J. P., G. F. Oster and J. C. Gerhart. 1986. Kinematics of gray crescent formation in *Xenopus* eggs: The displacement of subcortical cytoplasm relative to the egg surface. *Dev. Biol.* 113: 484–500.
- Vorwald-Denholtz, P. P. and E. M. De Robertis. 2011. Temporal pattern of the posterior expression of *Wingless* in *Drosophila* blastoderm. *Gene Expr. Patterns* 11: 456–463.
- Wacker, S. A., C. L. McNulty, A. J. Durston. 2004. The initiation of Hox gene expression in *Xenopus laevis* is controlled by Brachyury and BMP4. *Dev. Biol.* 266: 123–137.

Walentek, P., I. Schneider, A. Schweickert, and M. Blum. 2013. Wnt11b is involved in cilia-mediated symmetry breakage during *Xenopus* left-right development. *PLOS ONE* 8: e73646.

Warga, R. M. and C. B. Kimmel. 1990. Cell movements during epiboly and gastrulation in zebrafish. *Development* 108: 569–580.

Weaver C., G. H. Farr III, W. Pan, B. A. Rowning, J. Wang, J. Mao, D. Wu, L. Li, C. A. Larabell and D. Kimelman. 2003. GBP binds kinesin light chains and translocates during cortical rotation in *Xenopus* eggs. *Development* 130: 5425–5436.

Weaver, C. and D. Kimelman. 2004. Move it or lose it: Axis specification in *Xenopus*. *Development* 131: 3491–3499.

Wen, J. W. and R. Winklbauer. 2017. Ingression-type cell migration drives vegetal endoderm internalisation in the *Xenopus* gastrula. *Elife* 6: 27190.

Wessely, O., J. I. Kim, D. Geissert, U. Tran and E. M. De Robertis. 2004. Analysis of Spemann organizer formation in *Xenopus* embryos by cDNA macroarrays. *Dev. Biol.* 269: 552–566.

White, J. A. and J. Heasman. 2008. Maternal control of pattern formation in *Xenopus laevis*. *J. Exp. Zool. (MDE)* 310B: 73–84.

Wilson, P. A. and A. Hemmati-Brivanlou. 1995. Induction of epidermis and inhibition of neural fate by BMP-4. *Nature* 376: 331–333.

Wilson, P. A. and R. Keller. 1991. Cell rearrangement during gastrulation of *Xenopus*: Direct observation of cultured explants. *Development* 112: 289–300.

Winklbauer, R. and E. W. Damm. 2012. Internalizing the vegetal cell mass before and during amphibian gastrulation: Vegetal rotation and related movements. *Wiley Interdiscip. Dev. Biol.* 1: 301–306.

Winklbauer, R. and M. Schürfeld. 1999. Vegetal rotation, a new gastrulation movement involved in the internalization of the mesoderm and endoderm in *Xenopus*. *Development* 126: 3703–3713.

Xu, P. F., N. Houssin, K. F. Ferri-Lagneau, B. Thisse and C. Thisse. 2014. Construction of a vertebrate embryo from two opposing morphogen gradients. *Science* 344: 87–89.

Yang, J., C. Tan, R. S. Darken, P. A. Wilson and P. S. Klein. 2002. b-Catenin/Tcf-regulated transcription prior to the midblastula transition. *Development* 129: 5743–5752.

Yao, J. and D. S. Kessler. 2001. Goosecoid promotes head organizer activity by direct repression of Xwnt8 in Spemann's organizer. *Development* 128: 2975–2987.

Yost, C., M. Torres, J. R. Miller, E. Huang, D. Kimelman and R. T. Moon. 1996. The axis-inducing ability, stability, and subcellular localization of b-catenin are regulated in *Xenopus* embryos by glycogen synthase kinase-3. *Genes Dev.* 10: 1443–1454.

Zhang, J., D. W. Houston, M. L. King, C. Payne, C. Wylie and J. Heasman. 1998. The role of maternal VegT in establishing the primary germ layers in *Xenopus* embryos. *Cell* 94: 515–524.

Zimmerman, L. B., J. M. de Jesús-Escobar and R. M. Harland. 1996. The Spemann organizer signal noggin binds and inactivates bone morphogenesis protein 4. *Cell* 86: 599–606.