

Developmental Biology 13e, Chapter 19 Literature Cited

Aoyama, H. and K. Asamoto. 2000. The developmental fate of the rostral/caudal half of a somite for vertebra and rib formation: Experimental confirmation of the resegmentation theory using chick-quail chimeras. *Mech. Dev.* 99: 71–82.

Aszódi, A., D. Chan, E. Hunziker, J. F. Bateman and R. Fassler. 1998. Collagen II is essential for the removal of the notochord and the formation of intervertebral discs. *J. Cell Biol.* 143: 1399–1412.

Atit, R. and 7 others. 2006. Beta-catenin activation is necessary and sufficient to specify the dorsal dermal fate in the mouse. *Dev. Biol.* 296: 164–176.

Aulehla, A. and 7 others. 2008. A beta-catenin gradient links the clock and wavefront systems in mouse embryo segmentation. *Nat. Cell Biol.* 10: 186–193.

Aulehla, A., C. Wehrle, B. Brand-Saberi, R. Kemler, A. Gossler, B. Kanzler and B. G. Herrmann. 2003. Wnt3a plays a major role in the segmentation clock controlling somitogenesis. *Dev Cell.* 4: 395–406.

Barnes, G. L., P. G. Alexander, C. W. Hsu, B. D. Mariani and R. S. Tuan. 1997. Cloning and characterization of chicken paraxis: A regulator of paraxial mesoderm development and somite formation. *Dev. Biol.* 189: 95–111.

Barrios, A., R. J. Poole, L. Durbin, C. Brennan, N. Holder, and S. W. Wilson. 2003. Eph/Ephrin signaling regulates the mesenchymal-to-epithelial transition of the paraxial mesoderm during somite morphogenesis. *Curr. Biol.* 13: 1571–1582.

Beck C. W. 2015. Development of the vertebrate tailbud. *Wiley Interdiscip. Rev. Dev. Biol.* 4: 33–44.

Ben-Yair, R. and C. Kalcheim. 2005. Lineage analysis of the avian dermomyotome sheet reveals the existence of single cells with both dermal and muscle progenitor fates. *Development* 132: 689–701.

Bénazéraf, B. and O. Pourquié. 2013. Formation and segmentation of the vertebrate body axis. *Annu. Rev. Cell Dev. Biol.* 29: 1–26.

Bénazéraf, B., P. Francois, R. E. Baker, N. Denans, C. D. Little, and O. Pourquié. 2010. A random cell motility gradient downstream of FGF controls elongation of an amniote embryo. *Nature*. 466: 248–252.

Blair, H. C., A. J. Kahn, E. C. Crouch, J. J. Jeffrey and S. L. Teitelbaum. 1986. Isolated osteoclasts resorb the organic and inorganic components of bone. *J. Cell Biol.* 102: 1164–1172.

Blitz, E. and 9 others. 2009. Bone ridge patterning during musculoskeletal assembly is mediated through SCX regulation of Bmp4 at the tendon-skeleton junction. *Dev. Cell.* 17: 861–873.

Boettiger, D., M. Enomoto-Iwamoto, H. Y. Yoon, U. Hofer, A. S. Menko and R. Chiquet-Ehrismann. 1995. Regulation of integrin $\alpha 5 \beta 1$ affinity during myogenic differentiation. *Dev. Biol.* 169: 261–272.

Borycki, A., A. M. Brown and C. P. Emerson Jr. 2000. Shh and Wnt signaling pathways converge to control Gli gene activation in avian somites. *Development* 127: 2075–2087.

Bothe, I. and S. Dietrich. 2006. The molecular setup of the avian head mesoderm and its implication for craniofacial myogenesis. *Dev. Dyn.* 235: 2845–2860.

Bouldin, C. M., C. D. Snelson, G. H. Farr III, and D. Kimelman. 2014. Restricted expression of *cdc25a* in the tailbud is essential for formation of the zebrafish posterior body. *Genes Dev.* 28: 384–95.

Brand-Saberi, B., J. Wilting, C. Ebensperger and B. Christ. 1996. The formation of somite compartments in the avian embryo. *Int. J. Dev. Biol.* 40: 411–420.

Brent, A. E. and C. J. Tabin. 2004. FGF acts directly on the somitic tendon progenitors through the Ets transcription factors Pea3 and Erm to regulate scleraxis expression. *Development* 131: 3885–3896.

Brent, A. E., R. Schweitzer and C. J. Tabin. 2003. A somitic compartment of tendon precursors. *Cell* 113: 235–248.

Brent, A. E., T. Braun and C. J. Tabin. 2005. Genetic analysis of interactions between the somatic muscle, cartilage, and tendon cell lineages during mouse development. *Development* 132: 515–528.

Brill, G., N. Kahane, C. Carmeli, D. von Schack, Y. A. Barde and C. Kalcheim. 1995. Epithelial-mesenchymal conversion of dermatome progenitors requires neural tube-derived signals: Characterization of the role of Neurotrophin-3. *Development* 121: 2583–2594.

Bruder, S. P. and A. I. Caplan. 1989. Cellular and molecular events during embryonic bone development. *Connect. Tiss. Res.* 20: 65–71.

Buckingham, M., L. Bajard, P. Daubas, M. Esner, M. Lagha, F. Relaix and D. Rocancourt. 2006. Myogenic progenitor cells in the mouse embryo are marked by the expression of Pax3/7 genes that regulate their survival and myogenic potential. *Anat. Embryol.* 211: S51–S56.

Burgess, R., P. Cserjesi, K. L. Ligon and E. N. Olson. 1995. Paraxis: A basic helix-loop-helix protein expressed in paraxial mesoderm and developing somites. *Dev. Biol.* 168: 296–306.

Burke, A. C. and J. L. Nowicki. 2003. A new view of patterning domains in the vertebrate mesoderm. *Dev. Cell* 4: 159–165.

Carapuço, M., A. Nóvoa, N. Bobola and M. Mallo. 2005. Hox genes specify vertebral types in the presomitic mesoderm. *Genes Dev.* 19: 2116–2121.

Carron, C. and D. L. Shi. 2015 Specification of anteroposterior axis by combinatorial signaling during *Xenopus* development. *Wiley Interdiscip. Rev. Dev. Biol.* 5: 150–168.

Casaca, A., A. C. Santos, and M. Mallo. 2014. Controlling Hox gene expression and activity to build the vertebrate axial skeleton. *Dev. Dyn.* 243: 24–36.

Casaca, A., A. Nóvoa and M. Mallo. 2016. Hoxb6 can interfere with somitogenesis in the posterior embryo through a mechanism independent of its rib-promoting activity. *Development* 143: 437–448.

Cayuso, J., Q. Xu, and D. G. Wilkinson. 2015. Mechanisms of boundary formation by Eph receptor and ephrin signaling. *Dev. Biol.* 401: 122–131.

Chalamalasetty, R. B., R. J. Garriock, W. C. Dunty Jr., M. W. Kennedy, P. Jailwala, H. Si and T. P. Yamaguchi. 2014. Mesogenin 1 is a master regulator of paraxial presomitic mesoderm differentiation. *Development* 141: 4285–4297.

Chan, W. C., T. Y. Au, V. Tam, K. S. Cheah, and D. Chan. 2014. Coming together is a beginning: the making of an intervertebral disc. *Birth Defects Res. C Embryol. Today* 102: 83–100.

Chapman, D. L. and V. E. Papaioannou. 1998. Three neural tubes in mouse embryos with mutations in the T-box gene *Tbx6*. *Nature* 391: 695–697.

Chen, C.-M., N. Kraut, M. Groudine and H. Weintraub. 1996. I-mf, a novel myogenic repressor, interacts with members of the MyoD family. *Cell* 86: 731–741.

Cheung, T. H., N. L. Quach, G. W. Charville, L. Liu, L. Park, A. Edalati, B. Yoo, P. Hoang and T. A. Rando. 2012. Maintenance of muscle stem-cell quiescence by microRNA-489. *Nature* 482: 524–528.

Chipman, A. D. and M. Akam. 2008. The segmentation cascade in the centipede *Strigamia maritima*: Involvement of the Notch pathway and pair-rule gene homologues. *Dev. Biol.* 319: 160–169.

Choi, K. S. and B. D. Harfe. 2011. Hedgehog signaling is required for formation of the notochord sheath and patterning of nuclei pulposi within the intervertebral discs. *Proc. Natl. Acad. Sci. USA* 108: 9484–9489.

Choi, K. S., C. Lee, and B. D. Harfe. 2012. Sonic hedgehog in the notochord is sufficient for patterning of the intervertebral discs. *Mech. Dev.* 129: 255–262.

Choi, K. S., M. J. Cohn and B. D. Harfe. 2008. Identification of nucleus pulposus precursor cells and notochordal remnants in the mouse: implications for disk degeneration and chordoma formation. *Dev. Dyn.* 237: 3953–3958.

Christ, B. and C. P. Ohrdahl. 1995. Early stages of chick somite development. *Anat. Embryol.* 191: 381–396.

Christ, B., R. Huang and M. Scaal. 2007. Amniote somite derivatives. *Dev. Dyn.* 236: 2382–2396.

Cooke, J. and E. C. Zeeman. 1976. A clock and wavefront model for control of the number of repeated structures during animal morphogenesis. *J. Theor. Biol.* 58: 455–476.

Cooper, K. L., S. Oh, Y. Sung, R. R. Dasari, M. W. Kirschner and C. J. Tabin. 2013. Multiple phases of chondrocyte enlargement underlie differences in skeletal proportions. *Nature* 495: 375–378.

Corruccini, R. S. and C. L. Beecher. 1982. Occlusal variation related to soft diet in a nonhuman primate. *Science* 218: 74–76.

Corruccini, R. S. and C. L. Beecher. 1984. Occlusofacial morphological integration lowered in baboons raised on soft diet. *J. Craniofac. Genet. Dev. Biol.* 4: 135–142.

Cossu, G., R. Kelly, S. Tajbakhsh, S. Di Donna, E. Vivarelli and M. Buckingham. 1996a. Activation of different myogenic pathways: myf-5 is induced by the neural tube and MyoD by the dorsal ectoderm in mouse paraxial mesoderm. *Development* 122: 429–437.

Cossu, G., S. Tajbakhsh and M. Buckingham. 1996b. How is myogenesis initiated in the embryo? *Trends Genet.* 12: 218–223.

Crist, C. G., D. Montarras and M. Buckingham. 2012. Muscle satellite cells are primed for myogenesis but maintain quiescence with sequestration of Myf5 mRNA targeted by microRNA-31 in mRNP granules. *Cell Stem Cell.* 11: 118–126.

Cunningham, T. J. and 7 others. 2015. Retinoic acid activity in undifferentiated neural progenitors is sufficient to fulfill its role in restricting Fgf8 expression for somitogenesis. *PLoS ONE.* 10: e0137894.

Cunningham, T. J. and G. Duester. 2015. Mechanisms of retinoic acid signalling and its roles in organ and limb development. *Nat. Rev. Mol. Cell Biol.* 16: 110–123.

Dale, J. K., M. Maroto, M.-L. Dequeant, M. Malaper, M. McGrew and O. Pourquié. 2003. Periodic Notch inhibition by Lunatic Fringe underlies the chick segmentation clock. *Nature* 421: 275–278.

David, J. D., W. M. See and C. A. Higginbotham. 1981. Fusion of chick embryo skeletal myoblasts: Role of calcium influx preceding membrane union. *Dev. Biol.* 82: 297–307.

Delfini, M.-C., M. De La Celle, J. Gros, O. Serralbo, I. Marics, M. Seux, M. Scaal and C. Marcelle. 2009. The timing of emergence of muscle progenitors is controlled by an FGF/ERK/Snail1 pathway. *Dev. Biol.* 333: 229–237.

Denans, N., T. Iimura, and O. Pourquié. 2015. Hox genes control vertebrate body elongation by collinear Wnt repression. *Elife* 26: 4.

Denetclaw, W. F. and C. P. Ordahl. 2000. The growth of the dermomyotome and formation of early myotome lineages in thoracolumbar somites of chicken embryos. *Development* 127: 893–905.

Denkers, N., P. Garcia-Villalba, C. K. Rodesch, K. R. Nielson and T. J. Mauch. 2004. FISHing for chick genes: Triple-label whole mount fluorescence in situ hybridization detects simultaneous and overlapping gene expression in avian embryos. *Dev. Dyn.* 229: 651–657.

Dequéant, M. L. and O. Pourquié. 2008. Segmental patterning of the vertebrate embryonic axis. *Nat. Rev. Genet.* 9: 370–382.

Deries, M., A. B. Gonçalves, R. Vaz, G. G. Martins, G. Rodrigues and S. Thorsteinsdóttir. 2012. Extracellular matrix remodeling accompanies axial muscle development and morphogenesis in the mouse. *Dev. Dyn.* 241: 350–364.

Deries, M., R. Schweitzer and M. J. Duxson. 2010. Developmental fate of the mammalian myotome. *Dev. Dyn.* 239: 2898–2910.

Deschamps, J. and D. Duboule. 2017. Embryonic timing, axial stem cells, chromatin dynamics, and the Hox clock. *Genes Dev.* 3: 1406–1416.

Devoto, S. H. and 8 others. 2006. Generality of vertebrate developmental patterns: evidence for a dermomyotome in fish. *Evol. Dev.* 8: 101–110. Erratum in *Evol. Dev.* 2006. 2: 239.

Dietrich, S., F. R. Schubert, C. Healy, P. T. Sharpe and A. Lumsden. 1998. Specification of hypaxial musculature. *Development* 125: 2235–2249.

Doherty, K. R. and 7 others. 2005. Normal myoblast fusion requires myoferlin. *Development* 132: 5565–5575.

Dray, N, A. Lawton, A. Nandi, D. Jülich, T. Emonet, and S. A. Holley. 2013. Cell-fibronectin interactions propel vertebrate trunk elongation via tissue mechanics. *Curr. Biol.* 23: 1335–1341.

Duan, R. and P. J. Gallagher. 2009. Dependence of myoblast fusion on a cortical actin wall and nonmuscle myosin IIA. *Dev. Biol.* 325: 374–385.

Duboule, D. and G. Morata. 1994. Collinearity and functional hierarchy among genes of the homeotic complexes. *Trends Genet.* 10: 358–364.

Dubrulle, J. and O. Pourquié. 2004. Fgf8 mRNA decay establishes a gradient that couples axial elongation to patterning in the vertebrate embryo. *Nature* 427: 419–422.

Dubrulle, J., M. J. McGrew and O. Pourquié. 2001. FGF signaling controls somite boundary position and regulates segmentation clock control of spatiotemporal Hox gene activation. *Cell* 106: 219–232.

Dunty, W. C., Jr., K. K. Biris, R. B. Chalamalasetty, M. M. Taketo, M. Lewandoski, and T. P. Yamaguchi. 2008. Wnt3a/beta-catenin signaling controls posterior body development by coordinating mesoderm formation and segmentation. *Development* 135: 85–94.

Durbin, L. and 8 others. 1998. Eph signaling is required for segmentation and differentiation of the somites. *Genes Dev.* 12: 3096–3109.

Durland, J. L., M. Sferlazzo, M. Logan and A. C. Burke. 2008. Visualizing the lateral somitic frontier in the Prx1Cre transgenic mouse. *J. Anat.* 212: 590–602.

Eckalbar, W. L. and 12 others. 2012. Somitogenesis in the anole lizard and alligator reveals evolutionary convergence and divergence in the amniote segmentation clock. *Dev. Biol.* 363: 308–319.

Ellis, K., B. D. Hoffman, and M. Bagnat. 2013b. The vacuole within: How cellular organization dictates notochord function. *Bioarchitecture* 3: 64–68.

Ellis, K., J. Bagwell, and M. Bagnat. 2013a. Notochord vacuoles are lysosome-related organelles that function in axis and spine morphogenesis. *J. Cell Biol.* 200: 667–679.

Emery, A. E. 2002. The muscular dystrophies. *Lancet* 359: 687–695.

Evans, D. J. and D. M. Noden. 2006. Spatial relations between avian craniofacial neural crest and paraxial mesoderm cells. *Dev. Dyn.* 235: 1310–1325.

Fagotto, F., R. Winklbauer, and N. Rohani. 2014. Ephrin-Eph signaling in embryonic tissue separation. *Cell Adh. Migr.* 8: 308–326.

Fan, C. M. and M. Tessier-Lavigne. 1994. Patterning of mammalian somites by surface ectoderm and notochord: Evidence for sclerotome induction by a hedgehog homolog. *Cell* 79: 1175–1186.

Fisher, R. E., H. F. Smith, K. Kusumi, E. E. Tassone, A. Rawls and J. Wilson-Rawls. 2012. Mutations in the Notch pathway alter the patterning of multifidus. *Anat. Rec.* 295: 32–39.

Garriock, R. J., R. B. Chalamalasetty, M. W. Kennedy, L. C. Canizales, M. Lewandoski, and T. P. Yamaguchi. 2015. Lineage tracing of neuromesodermal progenitors reveals novel Wnt-dependent roles in trunk progenitor cell maintenance and differentiation. *Development* 142: 1628–1638.

Gerber, H. P., T. H. Vu, A. M. Ryan, J. Kowalski, Z. Werb and N. Ferrara. 1999. VEGF couples hypertrophic cartilage remodeling, ossification, and angiogenesis during endochondral bone formation. *Nature Med.* 5: 623–628.

Gerhart, J. and 9 others. 2011. Myo/Nog cell regulation of bone morphogenetic protein signaling in the blastocyst is essential for normal morphogenesis and striated muscle lineage specification. *Dev. Biol.* 359: 12–25.

Gerhart, J., J. Elder, C. Neely, J. Schure, T. Kvist, K. Knudsen and M. George-Weinstein. 2006. MyoD-positive epiblast cells regulate skeletal muscle differentiation in the embryo. *J. Cell Biol.* 175: 283–292.

Gomez, C., and O. Pourquié. 2009. Developmental control of segment numbers in vertebrates. *J. Exp. Zool.* 312B: 533–544.

Gomez, C., E. M. Ozbudak, J. Wunderlich, D. Baumann, J. Lewis, and O. Pourquié. 2008. Control of segment number in vertebrate embryos. *Nature* 454: 335–339.

Gray, R. S. and 7 others. 2014. Loss of col8a1a function during zebrafish embryogenesis results in congenital vertebral malformations. *Dev. Biol.* 386: 72–85.

Gros, J., M. Manceau, V. Thomé, and C. Marcelle. 2005. A common somitic origin for embryonic progenitors and satellite cells. *Nature* 435: 954–958.

Guehring, T., G. Wilde, M. Sumner, T. Grünhagen, G. B. Karney, U. K. Tirlapur and J. P. Urban. 2009. Notochordal intervertebral disc cells: Sensitivity to nutrient deprivation. *Arthritis Rheum.* 60: 1026–1034.

Guerreiro, I. and 8 others. 2013. Role of a polymorphism in a Hox/Pax-responsive enhancer in the evolution of the vertebrate spine. *Proc. Natl. Acad. Sci USA* 110: 10682–10686.

Gurung, R. and V. K. Parnaik. 2012. Cyclin D3 promotes myogenic differentiation and Pax7 transcription. *J. Cell Biochem.* 113: 209–219.

Haigh, J. J., H. P. Gerber, N. Ferrara and E. F. Wagner. 2000. Conditional inactivation of VEGF-A in areas of collagen 2a1 expression results in embryonic lethality in the heterozygous state. *Development* 127: 1445–1453.

Halata, Z., M. Grim and B. Christ. 1990. Origin of spinal cord meninges, sheaths of peripheral nerves, and cutaneous receptors including Merkel cells. An experimental and ultrastructural study with avian chimeras. *Anat. Embryol.* 182: 529–537.

Harel, I., E. Nathan, L. Tirosh-Finkel, H. Zigdon, N. Guimarães-Camboa, S. M. Evans and E. Tzahor. 2009. Distinct origins and genetic programs of head muscle satellite cells. *Dev. Cell* 16: 822–832.

Hatta, K., S. Takagi, H. Fujisawa and M. Takeichi. 1987. Spatial and temporal expression of N-cadherin cell adhesion molecule correlated with morphogenetic processes of chicken embryo. *Dev. Biol.* 120: 215–227.

Henrique, D., E. Abranches, L. Verrier, and K. G. Storey. 2015. Neuromesodermal progenitors and the making of the spinal cord. *Development* 142: 2864–2875.

Hirst, C. E. and C. Marcelle. 2015. The avian embryo as a model system for skeletal myogenesis. *Results Probl. Cell Differ.* 56: 99–122.

Ho, A. T. V., S. Hayashi, D. Bröhl, F. Auradé, R. Rattenbach and F. Relaix. 2011. Neural crest cell lineage restricts skeletal muscle progenitor cell differentiation through Neuregulin1-ErbB3 signaling. *Dev. Cell* 21: 273–287.

Holley, S. A. and C. Nüsslein-Volhard. 2000. Somitogenesis in zebrafish. *Curr. Top. Dev. Biol.* 47: 247–277.

Horsley, V., K. M. Jansen, S. T. Mills and G. K. Pavlath. 2003. IL-4 acts as a myoblast recruitment factor during mammalian muscle growth. *Cell* 113: 483–494.

Huang, R., Q. Zhi, B. Brand-Saberi, and B. Christ. 2000. New experimental evidence for somite resegmentation. *Anat. Embryol. (Berlin)* 202: 195–200.

Hubaud, A. and O. Pourquié. 2014. Signalling dynamics in vertebrate segmentation. *Nat. Rev. Mol. Cell Biol.* 15: 709–721.

Imura, T. and O. Pourquié. 2006. Collinear activation of *Hoxb* genes during gastrulation is linked to mesoderm cell ingression. *Nature* 442: 568–571.

Iulianella, A., B. Beckett, M. Petkovich, and D. Lohnes. 1999. A molecular basis for retinoic acid-induced axial truncation. *Dev. Biol.* 205: 33–48.

Izpisúa-Belmonte, J. C., H. Falkenstein, P. Dollé, A. Renucci, and D. Duboule. 1991a. Murine genes related to the *Drosophila AbdB* homeotic genes are sequentially expressed during development of the posterior part of the body. *EMBO J.* 10: 2279–2289.

Izpisúa-Belmonte, J.-C., C. Tickle, P. Dollé, L. Wolpert and D. Duboule. 1991b. Expression of the homeobox *Hox-4* genes and the specification of position in chick wing development. *Nature* 350: 585–589.

Jiang, Y. J., B. L. Aerne, L. Smithers, C. Haddon, D. Ish-Horowicz and J. Lewis. 2000. Notch signalling and the synchronization of the somite segmentation clock. *Nature* 408: 475–479.

Johnson, R. L., E. Laufer, R. D. Riddle and C. Tabin. 1994. Ectopic expression of Sonic hedgehog alters dorsal-ventral patterning of somites. *Cell* 79: 1165–1173.

Johnston, S. H., C. Rauskolb, R. Wilson, B. Prabhakaran, K.D. Irvine, T. F. Vogt. 1997. A family of mammalian Fringe genes implicated in boundary determination and the Notch pathway. *Development* 124: 2245–2254.

Jouve, C., I. Palmeirim, D. Henrique, J. Beckers, A. Gossler, D. Ish-Horowicz and O. Pourquié. 2000. Notch signalling is required for cyclic expression of the hairy-like gene HES1 in the presomitic mesoderm. *Development* 127: 1421–1429.

Jouve, C., T. Imura and O. Pourquié. 2002. Onset of segmentation clock in the chick embryo: Evidence for oscillations in the somite precursors in the primitive streak. *Development* 129: 1107–1117.

Jülich, D., A. P. Mould, E. Koper E, and S. A. Holley. 2009. Control of extracellular matrix assembly along tissue boundaries via Integrin and Eph/ Ephrin signaling. *Development* 136: 2913–2921.

Jülich, D., G. Cobb, A. M. Melo, P. McMillen, A. K. Lawton, S. G. Mochrie, E. Rhoades and S. A. Holley. 2015. Cross-scale integrin regulation organizes ECM and tissue topology. *Dev. Cell* 34: 33–44.

Kahn, A. J. and D. J. Simmons. 1975. Investigation of cell lineage in bone using a chimaera of chick and quail embryonic tissue. *Nature* 258: 325–327.

Kajimura, S., P. Seale, K. Kubota, E. Lunsford, J. V. Frangioni, S. P. Gygi and B. M. Spiegelman. 2009. Initiation of myoblast to brown fat switch by a PRDM-C/EBP- β transcriptional complex. *Nature* 460: 1154–1158.

Karsenty, G. and E. F. Wagner. 2002. Reaching a genetic and molecular understanding of skeletal development. *Dev. Cell* 2: 389–406.

Kato, N. and H. Aoyama. 1998. Dermomyotomal origin of the ribs as revealed by extirpation and transplantation experiments in chick and quail embryos. *Development* 125: 3437–3443.

Kieny, M., A. Mauger and P. Segel. 1972. Early regionalization of somitic mesoderm as studied by the development of the axial skeleton of the chick embryo. *Dev. Biol.* 28: 142–161.

Kimelman, D. and B. L. Martin. 2012. Anterior-posterior patterning in early development: Three strategies. *Wiley Interdiscip. Rev. Dev. Biol.* 1: 253–66.

Kimmel, C. B., W. W. Ballard, S. R. Kimmel, B. Ullmann, and T. F. Schilling. 1995. Stages of embryonic development of the zebrafish. *Dev. Dyn.* 203: 253–310.

Kmita, M. and D. Duboule. 2003. Organizing axes in time and space: 25 years of collinear tinkering. *Science* 301: 331–333.

Kondo, T. and D. Duboule. 1999. Breaking colinearity in the mouse HoxD complex. *Cell* 97: 407–417.

Koshida, S., Y. Kishimoto, H. Ustumi, T. Shimizu, M. Furutani-Seiki, H. Kondoh and S. Takada. 2005. Integrin α 5-dependent fibronectin accumulation for maintenance of somite boundaries in zebrafish embryos. *Dev. Cell* 8: 587–598.

Krol, A. J. and 8 others. 2011. Evolutionary plasticity of segmentation clock networks. *Development* 138: 2783–2792.

Kumar, S. and G. Duester. 2014. Retinoic acid controls body axis extension by directly repressing Fgf8 transcription. *Development* 141: 2972–2977.

Kuyyamudi, C., S. N. Menon and S. Sinha. 2021. Morphogen-regulated contact-mediated signaling between cells can drive the transitions underlying body segmentation in vertebrates. *Physical. Biol.* 19(1).

Lai, K. V. and 9 others. 2004. Conditional activation of *Akt* in adult skeletal muscle induces rapid hypertrophy. *J. Mol. Cell. Biol.* 24: 9295–9304.

Langman, J. 1981. *Medical Embryology*, 4th Ed. Williams & Wilkins, Baltimore, MD.

Lash, J. W. and K. M. Yamada. 1986. The adhesion recognition signal of fibronectin: A possible trigger mechanism for compaction during somitogenesis. In R. Bellairs, D. H. Ede and J. W. Lash (eds.), *Somites in Developing Embryos*. Plenum, New York, 201–208.

Lassar, A. B. and 6 others. 1989. MyoD is a sequence-specific DNA binding protein requiring a region of myc homology to bind to the muscle creatine kinase enhancer. *Cell* 58: 823–831.

Lawson, L. and B. D. Harfe. 2015. Notochord to nucleus pulposus transition. *Curr. Osteoporosis Rep.* 13: 336–41.

Lawton, A. K. and 7 others. 2013. Regulated tissue fluidity steers zebrafish body elongation. *Development* 140: 573–582.

Lee, S. J. 2004. Regulation of muscle mass by myostatin. *Annu. Rev. Cell Dev. Biol.* 20: 61–86.

Lewis, E. B. 1978. A gene complex controlling segmentation in *Drosophila*. *Nature* 276: 565–570.

Liang, Q. and 10 others. 2015. The roles of Mesp family proteins: functional diversity and redundancy in differentiation of pluripotent stem cells and mammalian mesodermal development. *Protein Cell* 6: 553–561.

Linask, K. K., C. Ludwig, M. D. Han, X. Liu, G. L. Radice and K. A. Knudsen. 1998. N-cadherin/catenin-mediated morphoregulation of somite formation. *Dev. Biol.* 202: 85–102.

Linde-Medina, M. and T. H. Smit. 2021. Molecular and mechanical cues for somite periodicity. *Front. Cell Dev. Biol.* 9: 753446.

Ling, B. M. and 10 others. 2012. Lysine methyltransferase G9a methylates the transcription factor MyoD and regulates skeletal muscle differentiation. *Proc. Natl. Acad. Sci. USA* 109: 841–846.

Long, F. 2012. Building strong bones: Molecular regulation of the osteoblast lineage. *Nat. Rev. Mol. Cell Biol.* 13: 27–38.

Maeda, T. and 12 others. 2011. Conversion of mechanical force into TGF- β -mediated biochemical signals. *Curr. Biol.* 21: 933–941.

Manolagas, S. C. and R. L. Jilka. 1995. Bone marrow, cytokines, and bone remodeling. *N. Engl. J. Med.* 332: 305–310.

Mansfield, J. H., E. Haller, N. D. Holland, and A. E. Brent. 2015. Development of somites and their derivatives in amphioxus, and implications for the evolution of vertebrate somites. *Evodevo* 6: 21.

Marcelle, C., M. R. Stark and M. Bronner-Fraser. 1997. Coordinate actions of BMPs, Wnts, Shh and Noggin mediate patterning of the dorsal somite. *Development* 124: 3955–3963.

Maroto, M., R. A. Bone, and J. K. Dale. 2012. Somitogenesis. *Development* 139: 2453–2456.

Martin, B. L. and D. Kimelman. 2012. Canonical Wnt signaling dynamically controls multiple stem cell fate decisions during vertebrate body formation. *Dev. Cell* 22: 223–232.

McCann, M. R., O. J. Tamplin, J. Rossant, and C. A. Seguin. 2011. Tracing notochord-derived cells using a Noto-cre mouse: implications for intervertebral disc development. *Dis. Model Mech.* 1: 73–82.

McGrew, M. J. and 9 others. 2008. Localised axial progenitor cell populations in the avian tailbud are not committed to a posterior Hox identity. *Development* 135: 2289–2299.

McMillen, P. and S. A. Holley. 2015. The tissue mechanics of vertebrate body elongation and segmentation. *Curr. Opin. Genet. Dev.* 32: 106–111.

McPherron, A. C., A. M. Lawler, and S. J. Lee. 1997. Regulation of skeletal muscle mass in mice by a new TGF-beta superfamily member. *Nature* 377: 83–90.

Meier, S. 1979. Development of the chick embryo mesoblast. Formation of the embryonic axis and establishment of the metameric pattern. *Dev. Biol.* 73: 25–45.

Menko, A. S. and D. Boettiger. 1987. Occupation of the extracellular matrix integrin is a control point for myogenic differentiation. *Cell* 51: 51–57.

Mittapalli, V. R., R. Huang, K. Patel, B. Christ and M. Scaal. 2005. Arthrotome: A specific joint forming compartment in the avian somite. *Dev. Dyn.* 234: 48–53.

Montavon, T. and D. Duboule. 2013. Chromatin organization and global regulation of Hox gene clusters. *Philos. Trans. R Soc. London B* 368: 20120367.

Morimoto, M., Y. Takahashi, M. Endo and Y. Saga. 2005. The Mesp2 transcription factor establishes segmental borders by suppressing Notch activity. *Nature* 435: 354–359.

Morin-Kensicki, E. M., E. Melancon and J. S. Eisen. 2002. Segmental relationship between somites and vertebral column in zebrafish. *Development* 129: 3851–3860.

Müller, F. and R. O’Rahilly. 1986. Somitic-vertebral correlation and vertebral levels in the human embryo. *Am. J. Anat.* 177: 3–19.

Münsterberg, A. E., J. Kitajewski, D. A. Bumcroft, A. P. McMahon and A. B. Lassar. 1995. Combinatorial signaling by sonic hedgehog and Wnt family members induce myogenic bHLH gene expression in the somite. *Genes Dev.* 9: 2911–2922.

Murray, P. J., F. A. Carrier and J. K. Dale. 2019. Cell cycle regulation of oscillations yields coupling of growth and form in a computational model of the presomitic mesoderm. *J. Theor. Biol.* 481: 75–83.

Naiche, L. A., N. Holder, and M. Lewandoski. 2011. FGF4 and FGF8 comprise the wavefront activity that controls somitogenesis. *Proc. Natl. Acad. Sci. USA* 108: 4018–4023.

Nakashima, K., X. Zhou, G. Kunkel, Z. Zhang, J. M. Deng, R. R. Behringer and B. de Crombrughe. 2002. The novel zinc finger-containing transcription factor osterix is required for osteoblast differentiation and bone formation. *Cell* 108: 17–29.

Nakaya, Y., S. Kuroda, Y. T. Katagiri, K. Kaibuchi and Y. Takahashi. 2004. Mesenchymal-epithelial transition during somitic segmentation is regulated by differential roles of Cdc42 and Rac1. *Dev. Cell* 7: 425–438.

Nameroff, M. and E. Munar. 1976. Inhibition of cellular differentiation by phospholipase C. II. Separation of fusion and recognition among myogenic cells. *Dev. Biol.* 49: 288–293.

Neijts, R., S. Simmini, F. Giuliani, C. van Rooijen, and J. Deschamps. 2014. Region-specific regulation of posterior axial elongation during vertebrate embryogenesis. *Dev. Dyn.* 243: 88–98.

Nguyen, P. D. and 14 others. 2014. Haematopoietic stem cell induction by somite-derived endothelial cells controlled by *meox1*. *Nature* 512: 314–318.

Nikaido, M., A. Kawakami, A. Sawada, M. Furutani-Seiki, H. Takeda, and K. Araki. 2002. Tbx24, encoding a T-box protein, is mutated in the zebrafish somite-segmentation mutant fused somites. *Nat. Genet.* 31: 195–199.

Nimmagadda, S., P. Geetha-Loganathan, M. Scaal, B. Christ and R. Huang. 2007. FGFs, Wnts and BMPs mediate induction of VEGFR-2 (Quek-1) expression during avian somite development. *Dev. Biol.* 305: 421–429.

Niwa, Y., H. Shimojo, A. Isomura, A. González, H. Miyachi and R. Kageyama. 2011. Different types of oscillations in Notch and FGF signaling regulate the spatiotemporal periodicity of somitogenesis. *Genes Dev.* 25: 1115–1120.

Noordermeer, D., M. Leleu, P. Schorderet, E. Joye, F. Chabaud, and D. Duboule. 2014. Temporal dynamics and developmental memory of 3D chromatin architecture at Hox gene loci. *Elife* 3: e02557.

Nowicki, J. L. and A. C. Burke. 2000. Hox genes and morphological identity: Axial versus lateral patterning in the vertebrate mesoderm. *Development* 127: 4265–4275.

Nowicki, J. L., R. Takimoto and A. C. Burke. 2003. The lateral somitic frontier: Dorso-ventral aspects of anterior-posterior regionalization in the embryo. *Mech. Dev.* 120: 227–240.

Nowotschin, S., A. Ferrer-Vaquer, D. Concepcion, V. E. Papaioannou, and A. K. Hadjantonakis. 2012. Interaction of Wnt3a, Msgn1 and Tbx6 in neural versus paraxial mesoderm lineage commitment and paraxial mesoderm differentiation in the mouse embryo. *Dev. Biol.* 367: 1–14.

Oberlender, S. A. and R. S. Tuan. 1994. Expression and functional involvement of N-cadherin in embryonic limb chondrogenesis. *Development* 120: 177–187.

Ohata, E., R. Tadokoro, Y. Sato, D. Saito and Y. Takahashi. 2009. Notch signal is sufficient to direct an endothelial conversion from non-endothelial somitic cells conveyed to the aortic region by CXCR4. *Dev. Biol.* 335: 33–42.

Olivera-Martinez, I., S. Missier, S. Fraboulet, J. Thélou and D. Dhouailly. 2002. Differential regulation of the chick dorsal thoracic dermal progenitors from medial dermamyotome. *Development.* 129: 4763–4772.

Ontell, M., D. Hughes and D. Bourke. 1988. Morphometric analysis of the developing mouse soleus muscle. *Am. J. Anat.* 181: 279–288.

Ordahl, C. P. 1993. Myogenic lineages within the developing somite. In M. Bernfield (ed.), *Molecular Basis of Morphogenesis*. Wiley-Liss, New York, 165–170.

Ordahl, C. P. and N. Le Douarin. 1992. Two myogenic lineages within the developing somite. *Development* 114: 339–353.

Ordahl, C. P., E. Berdugo, S. J. Venters, and W. F. J. Denetclaw. 2001. The dermomyotome dorsomedial lip drives growth and morphogenesis of both the primary myotome and dermomyotome epithelium. *Development* 128: 1731–1744.

Palmeirim, I., D. Henrique, D. Ish-Horowicz and O. Pourquié. 1997. Avian hairy gene expression identifies a molecular clock linked to vertebrate segmentation and somitogenesis. *Cell* 91: 639–648.

Pardanaud, L., D. Luon, M. Prigent, L.-M., Bourcheix, M. Catala and F. Dieterlen-Lievre 1996. Two distinct endothelial lineages in ontogeny, one of them related to hemopoiesis. *Development* 122: 1363–1371.

Peres J., A. McNulty, and A. Durston. 2006. Interaction between X-Delta-2 and Hox genes regulates segmentation and patterning of the anteroposterior axis. *Mech. Dev.* 123: 321–333.

Piette, J., J.-L. Bessereau, M. Huchet and J.-P. Changeux. 1990. Two adjacent MyoD1-binding sites regulate expression of the acetylcholine receptor α -subunit gene. *Nature* 345: 353–355.

Pourquié, O. 2011. Vertebrate segmentation: From cyclic gene networks to scoliosis. *Cell* 145: 650–663.

Pourquié, O. and P. P. Tam. 2001. A nomenclature for prospective somites and phases of cyclic gene expression in the presomitic mesoderm. *Dev. Cell.* 1: 619–620.

Pourquié, O., C. M Fan, M. Coltey, E. Hirsinger, Y. Watanabe, C. Bréant, P. Francis-West, P. Brickell, M. Tessier-Lavigne and N. M. Le Douarin. 1996. Lateral and axial signals involved in somite patterning: A role for BMP4. *Cell* 84: 461–471.

Pueyo, J. I., R. Lanfear and J. P. Couso. 2008. Ancestral Notch-mediated segmentation revealed in the cockroach *Periplaneta americana*. *Proc. Natl. Acad. Sci. USA* 105: 16614–16619.

Punch, V. G., A. E. Jones and M. A. Rudnicki. 2009. Transcriptional networks that regulate muscle stem cell function. *Wiley Interdiscip. Rev. Syst. Biol. Med.* 1: 128–140.

Relaix, F., D. Rocancourt, A. Mansouri and. Buckingham. 2005. A Pax3/Pax7-dependent population of skeletal muscle progenitor cells. *Nature* 435: 948–953.

Remak, R. 1850. *Untersuchungen über die Entwicklung der Wirbeltiere*. Remier, Berlin.

Reshef, R., M. Maroto and A. B. Lassar. 1998. Regulation of dorsal somitic cell fates: BMPs and Noggin control the timing and pattern of myogenic regulator expression. *Genes Dev.* 12: 290–303.

Richardson, B. E., S. J. Nowak and M. K. Baylies. 2008. Myoblast fusion in fly and vertebrates: New genes, new processes and new perspectives. *Traffic* 9: 1050–1059.

Rios, A. C., O. Serralbo, D. Salgado, and C. Marcelle. 2011. Neural crest regulates myogenesis through the transient activation of Notch. *Nature* 473: 532–535.

Risbud, M. V. and I. M. Shapiro. 2011. Notochordal cells in the adult intervertebral disc: New perspective on an old question. *Crit. Rev. Eukaryot. Gene Expr.* 21: 29–41.

Saga, Y. 2007. Segmental border is defined by the key transcription factor Mesp2, by means of the suppression of Notch activity. *Dev. Dyn.* 236: 1450–1455.

Saga, Y., N. Hata, H. Koseki, and M. M. Taketo. 1997. Mesp2: A novel mouse gene expressed in the presegmented mesoderm and essential for segmentation initiation. *Genes Dev.* 11: 1827–1839.

Sartorelli, V. and A. H. Juan. 2011. Sculpting chromatin beyond the double helix: Epigenetic control of skeletal myogenesis. *Curr. Top. Dev. Biol.* 96: 57–83.

Sato, Y., K. Yasuda and Y. Takahashi. 2002. Morphological boundary forms by a novel inductive event mediated by Lunatic-Fringe and Notch during somitic segmentation. *Development* 129: 3633–3644.

Sato, Y., T. Watanabe, D. Saito, T. Takahashi, S. Yoshida, J. Kohyama, E. Ohata, H. Okano and Y. Takahashi. 2008. Notch mediates the segmental specification of angioblasts in somites and their directed migration toward the dorsal aorta in avian embryos. *Dev. Cell* 14: 890–901.

Scaal, M. 2015. Early development of the vertebral column. *Semin. Cell Dev. Biol.* 49: 83–91.

Schweitzer, R. and 7 others. 2001. Analysis of the tendon cell fate using Scleraxis, a specific marker for tendons and ligaments. *Development* 128: 3855–3866.

Serralbo, O. and C. Marcelle. 2014. Migrating cells mediate long-range WNT signaling. *Development*. 141: 2057–2063.

Shefer, G., G. Rauner, Z. Yablonka-Reuveni and D. Benayahu. 2010. Reduced satellite cell numbers and myogenic capacity in aging can be alleviated by endurance exercise. *PLOS ONE* 5: e13307.

Smith, C. A. and R. S. Tuan. 1996. Functional involvement of Pax-1 in somite development: Somite dysmorphogenesis in chick embryos treated with Pax-1 paired-box antisense oligonucleotide. *Teratology* 52: 333–345.

Soroldoni, D., D. J. Jörg, L. G. Morelli, D. L. Richmond, J. Schindelin, F. Jülicher and A. C. Oates. 2014. Genetic oscillations: A Doppler effect in embryonic pattern formation. *Science* 345: 222–225.

Soshnikova, N. and D. Duboule. 2009. Epigenetic temporal control of mouse Hox genes in vivo. *Science* 324: 1320–1323.

Sosic, D., B. Brand-Saberi, C. Schmidt, B. Christ and E. N. Olson. 1997. Regulation of paraxis expression and somite formation by ectoderm- and neural tube-derived signals. *Dev. Biol.* 185: 229–243.

Sparrow, D. B. and 9 others. 2006. Mutation of the *LUNATIC FRINGE* gene in humans causes spondylocostal dysostosis with a severe vertebral phenotype. *Am. J. Hum. Genet.* 78: 28–37.

St.-Jacques, B., M. Hammerschmidt and A. P. McMahon. 1999. Indian hedgehog signaling regulates proliferation and differentiation of chondrocytes and is essential for bone formation. *Genes Dev.* 13: 2072–2086.

Stern, H. M., A. M. C. Brown and S. D. Hauschka. 1995. Myogenesis in paraxial mesoderm: Preferential induction by dorsal neural tube and by cells expressing *Wnt-1*. *Development* 121: 3675–3686.

Takahashi, Y., A.-H. Monsoro-Burg, M. Bontoux and N. M. Le Douarin. 1992. A role for Quox8 in the establishment of the dorsoventral pattern during vertebrate development. *Proc. Natl. Acad. Sci. USA* 89: 10237–10241.

Takahashi, Y., K. Koizumi, A. Takagi, S. Kitajima, T. Inoue, H. Koseki and Y. Saga. 2000. *Mesp2* initiates somite segmentation through the Notch signalling pathway. *Nat. Genet.* 25: 390–396.

Takahashi, Y., S. Kitajima, T. Inoue, J. Kanno, and Y. Saga. 2005. Differential contributions of *Mesp1* and *Mesp2* to the epithelialization and rostro-caudal patterning of somites. *Development* 132: 787–796.

Takemoto, T., M. Uchikawa, M. Yoshida, D. M. Bell, R. Lovell-Badge, V. E. Papaioannou and H. Kondoh. 2011. *Tbx6*-dependent *Sox2* regulation determines neural or mesodermal fate in axial stem cells. *Nature* 470: 394–398.

Tam, P. P. 1981. The control of somitogenesis in mouse embryos. *J. Embryol. Exp. Morphol.* 65: 103–128.

Tani, S., U.-I. Chung, S. Ohba and H. Hojo. 2020. Understanding paraxial mesoderm development and sclerotome specification for skeletal repair. *Exp. Mol. Med* 52, 1166–1177.

Tonegawa, A. and Y. Takahashi. 1998. Somitogenesis controlled by *Noggin*. *Dev. Biol.* 202: 172–182.

Tonegawa, A., N. Funayama, N. Ueno and Y. Takahashi. 1997. Mesodermal subdivision along the mediolateral axis in chicken controlled by different concentrations of BMP-4. *Development* 124: 1975–1984.

Tseng, Y. H. and 14 others. 2008. New role of bone morphogenetic protein 7 in brown adipogenesis and energy expenditure. *Nature* 454: 1000–1004.

Tzouanacou, E., A. Wegener, F. J. Wymeersch, V. Wilson, and J. F. Nicolas. 2009. Redefining the progression of lineage segregations during mammalian embryogenesis by clonal analysis.

Dev. Cell. 17: 365–376.

van Eeden, F. J., S. A. Holley, P. Haffter, and C. Nüsslein-Volhard. 1998. Zebrafish segmentation and pair-rule patterning. *Dev. Genet.* 23: 65–76.

Varrela, J. 1992. Dimensional variation of craniofacial structures in relation to changing masticatory-functional demands. *Eur. J. Orthod.* 14: 31–36.

Vinagre, T., N. Moncaut, M. Carapucol, A. Novoal and J. Bom. 2010. Evidence for a myotomal Hox/Myf cascade governing nonautonomous control of rib specification within global vertebral domains. *Dev. Cell* 18: 655–661.

Wahi, K., M. S. Bochter, and S. E. Cole. 2014. The many roles of Notch signaling during vertebrate somitogenesis. *Semin. Cell Dev. Biol.* 49: 68–75.

Wang, Y. and R. Serra. 2012. PDGF mediates TGFb-induced migration during development of the spinous process. *Dev. Biol.* 365: 110–117.

Watanabe, T. and Y. Takahashi. 2010. Tissue morphogenesis coupled with cell shape changes. *Curr. Opin. Genet. Dev.* 20: 443–447.

Watanabe, T., Y. Sato, D. Saito, R. Tadokoro, and Y. Takahashi. 2009. EphrinB2 coordinates the formation of a morphological boundary and cell epithelialization during somite segmentation. *Proc. Natl. Acad. Sci. USA* 106: 7467–7472.

Wellik, D. M. and M. R. Capecchi. 2003. *Hox10* and *Hox11* genes are required to globally pattern the mammalian skeleton. *Science* 301: 363–367.

Wilm, B., R. G. James, T. M. Schultheiss and B. L. M. Hogan. 2004. The forkhead genes, *Foxc1* and *Foxc2*, regulate paraxial versus intermediate mesoderm cell fate. *Dev. Biol.* 271: 176–189.

Windner, S. E., N. C. Bird, S. E. Patterson, R. A. Doris, and S. H. Devoto. 2012. Fss/Tbx6 is required for central dermomyotome cell fate in zebrafish. *Biol. Open.* 1: 806–814.

Winslow, B. B., R. Takimoto-Kimura and A. C. Burke. 2007. Global patterning of the vertebrate mesoderm. *Dev. Dyn.* 236: 2371–2381.

- Yabe, T. and S. Takada. 2012. Mesogenin causes embryonic mesoderm progenitors to differentiate during development of zebrafish tail somites. *Dev. Biol.* 370: 213–222.
- Yamashita, S. and 9 others. 2012. L-Sox5 and Sox6 proteins enhance chondrogenic miR-140 microRNA expression by strengthening dimeric Sox9 activity. *J. Biol. Chem.* 287: 22206–22215.
- Yang, S. Y., M. Alnaqeeb, H. Simpson and G. Goldspink. 1996. Cloning and characterization of an IGF-I isoform expressed in skeletal muscle subjected to stretch. *J. Musc. Res. Cell Motil.* 17: 487–495.
- Yin, M., C. Gentili, E. Koyama, M. Zasloff and M. Pacifici. 2002. Antiangiogenic treatment delays chondrocyte maturation and bone formation during limb skeletogenesis. *J. Bone Miner. Res.* 17: 56–65.
- Young, T. and 12 others. 2009. Cdx and Hox genes differentially regulate posterior axial growth in mammalian embryos. *Dev. Cell* 17: 516–526.
- Zákány, J., M. Kmita, P. Alarcon, J.-L. de la Pompa and D. Duboule. 2001. Localized and transient transcription of Hox genes suggests a link between patterning and the segmentation clock. *Cell* 106: 207–217.
- Zhang, N., C. R. Norton, and T. Gridley. 2002. Segmentation defects of Notch pathway mutants and absence of a synergistic phenotype in *lunatic fringe/radical fringe* double mutant mice. *Genesis* 33: 21–28.